This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Chronic probiotic supplementation with or without glutamine does not influence the eHsp72 response to a multi-day ultra-endurance exercise event

PLEASE CITE THE PUBLISHED VERSION

<https://doi.org/10.1139/apnm-2017-0131>

PUBLISHER

NRC Research Press (© the Authors)

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Marshall, Hannah, Bryna C.R. Christmas, Craig Anthony Suckling, Justin D. Roberts, Josh Foster, and Lee Taylor. 2017. "Chronic Probiotic Supplementation with or Without Glutamine Does Not Influence the Ehsp72 Response to a Multi-day Ultra-endurance Exercise Event". figshare. <https://hdl.handle.net/2134/28352>.

Applied Physiology, Nutrition, and Metabolism
Physiologie appliquée, nutrition et métabolisme

**Chronic probiotic supplementation with or without
glutamine does not influence the eHsp72 response to a
multi-day ultra-endurance exercise event.**

Journal:	<i>Applied Physiology, Nutrition, and Metabolism</i>
Manuscript ID	apnm-2017-0131.R1
Manuscript Type:	Article
Date Submitted by the Author:	07-Apr-2017
Complete List of Authors:	Marshall, Hannah; University of Bedfordshire, Institute of Sport and Physical Activity Research (ISPAR) Christmas, Bryna; Qatar University, Sport Science Program, College of Arts and Sciences; Suckling, Craig; Anglia Ruskin University Roberts, Justin; Anglia Ruskin University Foster, Josh; University of Bedfordshire, Institute of Sport and Physical Activity Research (ISPAR) Taylor, Lee; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Athlete Health and Performance Research Centre
Keyword:	extracellular heat shock protein 72, probiotics, Ultra-endurance, supplementation, exercise nutrition < exercise

SCHOLARONE™
Manuscripts

1 **Chronic probiotic supplementation with or without glutamine does not influence**
2 **the eHsp72 response to a multi-day ultra-endurance exercise event.**

3

4 **Hannah Marshall¹, Bryna Catherine Rose Christmas², Craig Andrew Suckling³, Justin D.**
5 **Roberts³, Josh Foster¹, Lee Taylor^{4*,5}**

6

7 ¹ Institute of Sport and Physical Activity Research (ISPAR), University of Bedfordshire, Bedford, UK.

8 ²Sport Science Program, College of Arts and Sciences, Qatar University, Doha, Qatar.

9 ³Cambridge Centre for Sport and Exercise Sciences, Department of Life Sciences, Anglia Ruskin
10 University, Cambridge, UK.

11 ⁴ASPETAR, Athlete Health and Performance Research Centre, Qatar Orthopaedic and Sports Medicine
12 Hospital, Aspire Zone, Doha, Qatar.

13 ⁵School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK.

14

15 **Corresponding Author:** Dr. Lee Taylor, ASPETAR, Athlete Health and Performance Research
16 Centre, Qatar Orthopaedic and Sports Medicine Hospital, Aspire Zone, Doha, Qatar. Email:
17 Lee.Taylor@aspetar.com. Phone: + (974) 4413 2403.

18

19

20

21

22

23

24

25

26

27

28

29

30

Abstract

Probiotic and glutamine supplementation increases tissue Hsp72, but their influence on extracellular Hsp72 (eHsp72) has not been investigated. The aim of this study was to investigate the effect of chronic probiotic supplementation, with or without glutamine, on eHsp72 concentration before and after an ultramarathon. Thirty-two participants were split into three independent groups, where they ingested probiotic capsules (PRO, $n = 11$), probiotic + glutamine powder (PGLn, $n = 10$) or no supplementation (CON, $n = 13$), over a 12-week period prior to commencement of the Marathon des Sables (MDS). eHsp72 concentration in the plasma was measured at baseline, 7 days pre-race, 6-8 hours post-race, and 7 days post-race. The MDS increased eHsp72 concentrations by 124% ($F_{1,3} = 22.716$, $p < 0.001$), but there was no difference in the response between groups. Additionally, PRO or PGLn supplementation did not modify pre- or post-MDS eHsp72 concentrations compared to CON ($p > 0.05$). In conclusion, The MDS caused a substantial increase in eHsp72 concentration indicating high levels of systemic stress. However, chronic PRO or PGLn supplementation did not effect eHsp72 compared to control pre- or post-MDS. Given the role of eHsp72 in immune activation, the commercially available supplements used in this study are unlikely to influence this cascade.

Key words: Ultra-endurance; extracellular heat shock protein 72; probiotics; desert; race; event.

61 **Introduction**

62

63 Exercise in extreme heat elicits physiological stress upon the human body (Nybo et al. 2001) which can
64 decrement physical performance (Nybo et al. 2014) and cognitive function (Taylor et al. 2015). In
65 addition, strenuous exercise coupled with high environmental temperatures (exercise heat-stress; EHS)
66 is a major risk factor for the development of exertional heat illnesses [EHI; (Armstrong et al. 2007)]. A
67 plethora of physiological responses including elevated core temperature (Binkley et al. 2002),
68 increased heart rate (HR) and a redistribution of blood flow (González-Alonso et al. 2008) are seen in
69 response to EHS (Nybo 2008; 2014). These responses initially act to protect the body from tissue
70 damage and protein denaturation during EHS, yet they can prove catastrophic if the total stress exceeds
71 a highly individualised threshold of tolerance [EHI; (Adams et al. 2015; Goforth et al. 2015;
72 Wijerathne et al. 2016)]. Nonetheless, humans willingly subject themselves to prolonged EHS when
73 completing endurance events such as the Marathon des Sables (MDS). The MDS is a 294.4 km 7 day
74 ultra-marathon completed across the Southern Moroccan portion of Sahara Desert. Ultra-endurance
75 events such as the MDS are becoming increasingly popular, with a substantial rise in competitor
76 numbers in recent years (Knechtle et al. 2011; da Fonseca-Engelhardt et al. 2013).

77

78 Exposure to EHS can increase extracellular heat shock protein (eHsp72) concentration (Walsh et al.
79 2001; Whitham et al. 2007; Sandström et al. 2008; Magalhaes et al. 2010; Périard et al. 2012; Gibson et
80 al. 2014), a protein which is thought to contribute to the immune response to EHS (Mizzen et al. 1988;
81 Asea 2005). This increase in eHsp72 is thought to prepare the immune system (immuno-stimulatory)
82 for subsequent EHS mediated homeostatic perturbations (Johnson et al. 2006). Although discussion
83 regarding the precise role of eHsp72 remain (Johnson and Fleshner 2006; Whitham et al. 2008),
84 evidence suggests that eHsp72 enhances the production of pro-inflammatory cytokines (Asea 2005)
85 and chemokines (Lehner et al. 2000). This implies the importance of eHsp72 during EHS, given the
86 role of pro-inflammatory cytokines in the pathophysiology of heat stroke (Lim et al. 2006; Leon et al.
87 2010). Previous *in vitro* (Wischmeyer 2002; Petrof et al. 2004; Tao et al. 2006) and *in vivo*
88 (Wischmeyer et al. 2001) research has demonstrated that glutamine increases intracellular Hsp72
89 [iHsp72 (epithelial cells)] concentrations, thus it is plausible to suggest that such interventions may
90 elevate eHsp72 concentrations. Because the release mechanism of Hsp72 into the extracellular

environment is still not fully defined, it is unknown if interventions that increase iHsp72 also influence eHsp72 concentrations.

Previous research has demonstrated exercise duration to be a key modulator of eHsp72 release (Febbraio et al. 2002; Marshall et al. 2006; Amorim et al. 2008), and therefore ultra-endurance events, defined as lasting > 6 h (Zaryski et al. 2005), are likely to induce a significant increase in eHsp72 concentration. The MDS (294.4 km, 7 days, 20-50°C) is physiologically challenging given that competitors must carry food and survival equipment for the duration of the event, with only rationed water provided each day, subsequently placing competitors at an increased risk of EHS and potentially EHI throughout the race. Therefore, a nutritional intervention which increases eHsp72 concentration prior to a multi-day ultra-endurance event, such as the MDS, could alter the associated immune response patterns in competitors. Whether or not a basal increase in eHsp72 is beneficial or detrimental in this regard is currently unknown. The additions of probiotics and glutamine have been shown to induce an increase in Hsp72 concentration in various cultured cell types (Wischmeyer 2002; Petrof et al. 2004; Tao et al. 2006). However, to our knowledge, no research has yet been conducted in relation to probiotic supplementation and the eHsp72 response to EHS relative to a multi-day ultra-endurance event with logistical and nutritional challenges, such as the MDS. Owing to its immuno-stimulatory actions, a probiotic or glutamine mediated increase in pre-race eHsp72 may influence the immune response pattern to EHS (MDS). This knowledge is important because any alteration in pro/anti-inflammatory responses to EHS could influence the severity of EHI (Lim and Mackinnon 2006). Despite previous studies demonstrating a significant increase in iHsp72 through probiotic and glutamine supplementation (Wischmeyer 2002; Petrof et al. 2004), no research has investigated if this alters eHsp72 concentrations prior to and following EHS.

This preliminary study aimed to investigate the effects of chronic (12-weeks) probiotic supplementation with (PGLn) or without (PRO) glutamine (both commercially available), on the eHsp72 response pre- and post-MDS completion. Owing to the effects of probiotics and glutamine on Hsp72 synthesis, it was hypothesised that PRO and PGLn would increase eHsp72 concentrations pre-MDS in comparison to a control group (CON). Additionally, it was hypothesised that the nutritional

120 interventions would reduce the stress and thus the inflammatory response to exercise, marked by a
121 blunted post-MDS eHsp72 concentration when compared to CON.

122

123 **Materials and methods**

124

125 **Participants**

126

127 Male and female entrants (aged 18-60 years) were contacted via email through the UK organiser
128 (RunUltra UK). Thirty-two MDS 2015 competitors (6 female, 24 male, age 41; range 23-53 years,
129 height 1.75 ± 0.08 m, body mass 76.89 ± 2.04 kg) met the inclusion criteria and were recruited for this
130 study. Participants were excluded from taking part if: (1) unable to attend all testing sessions; (2)
131 outside of desired age range (18 – 60 years); (3) unsatisfactory health-screen questionnaire completion
132 (i.e. potential history of heart abnormalities, hypertension, heart disease or diabetes); (4) suffering from
133 any musculo-skeletal injury that may have impaired participation in the study and/or endurance
134 training; (5) any known blood related disorder; (6) were taking over-the-counter and/or prescribed
135 medication which may have influenced normal participation (excluding inhalers for exercise-induced
136 asthma or the contraceptive pill); (7) were consuming any other commercial supplementation which
137 conflicted the study parameters (i.e. an alternative probiotic) – and were unwilling to refrain from use
138 of supplementation over the study period; and (8) any participant who was, or had the potential to
139 become, pregnant during the study. All participants provided written informed consent, were deemed
140 healthy and able to take part, and verbally confirmed that they were not currently undertaking any
141 probiotic or glutamine supplementation regime. Participants were instructed to avoid heat
142 acclimation/acclimatisation training sessions in the 48 h prior to all data collection visits (excluding the
143 immediate post-race sample). Of the 32 recruited participants, 94% were compliant to this requirement,
144 with one participant in both the PRO and CON groups undertaking a bout of heat acclimation training
145 48 h prior to data collection visit 2. Between data collection visit 2 and departure for the MDS, 31.25%
146 ($n = 10$) of participants undertook at least one heat training session [PRO: 2 sessions ($n = 5$), 3
147 sessions ($n = 1$); PGLn: 3 sessions ($n = 1$); CON: 2 sessions ($n = 2$), 4 sessions ($n = 1$)]. All
148 procedures were approved by the Anglia Ruskin University Ethics Committee, and conformed to the
149 Declaration of Helsinki.

150

151 Study overview

152

153 Following a randomised independent measures design; participants were assigned to one of three
154 experimental conditions utilizing commercially available dietary supplements:

155

156 *Probiotic capsules (PRO; n = 11)*: Participants were required to consume one capsule (Bio-
157 Acidophilus Forte, Biocare Ltd., Birmingham, UK) per day for the duration of the 12-week
158 intervention period. Each multi-strain capsule contained 150 mg.d⁻¹ *Lactobacillus acidophilus* (10
159 billion CFU.d⁻¹, *Lactobacillus acidophilus* CUL-60 [NCIMB 30157] and 10 billion CFU.d⁻¹
160 *Lactobacillus acidophilus* CUL-21 [NCIMB 30156]), 16.8 mg.d⁻¹ *Bifidobacterium bifidum and lactis*
161 (9.5 billion CFU.d⁻¹, *Bifidobacterium bifidum* CUL-20 [NCIMB 30172] and 0.5 billion CFU.d⁻¹
162 *Bifidobacterium animalis subspecies lactis* CUL-34 [NCIMB 30153]), and 55.8 mg.d⁻¹
163 *fructooligosaccharides*.

164

165 Probiotic + glutamine powder (*PGLn; n = 10*): Participants were required to consume 5 g powder (GI
166 Complex, UK) per day, mixed well in water or food. 2 billion, *Lactobacillus acidophilus* CUL-60
167 (NCIMB 30157), 2 billion; *Lactobacillus acidophilus* CUL-21 (NCIMB 30156); 50 million,
168 *Bifidobacterium bifidum* CUL-20 (NCIMB 30172); 0.95 billion, *Bifidobacterium animalis* subsp.
169 *Lactis* CUL-34 (NCIMB 30153); 5 billion, *Lactobacillus salivarius* CUL61 (NCIMB 30211), each 5 g
170 dose also contained 0.9 g L-Glutamine.

171

172 *Control (CON; n = 9)*: maintenance of regular diet.

173

174 Apparent daily adherence to intervention and control was self-reported at 100% for all groups. Due to
175 the nature of the study, blinding of groups was not possible, however, the PGLn group were unaware of
176 the addition of glutamine to their probiotic supplement.

177

178 Data was collected at four time points across the duration of the study, which consisted of three
179 laboratory visits (visits 1, 2, and 4, Anglia Ruskin University, Cambridge) and one field-based data
180 collection point (visit 3, Ouarzazate, Morocco), as detailed below (Fig. 1a).

181

182 Visit 1

183

184 This baseline data collection took place 12-weeks prior to the MDS. Participants arrived at the
185 environmentally controlled laboratory (18°C, 35% RH) in a fasted state (minimum 4 h fasted), with
186 confounding variables of alcohol (Taylor et al. 2010a), caffeine (Lu et al. 2008), generic
187 supplementation (Hillman et al. 2011) and smoking (Anbarasi et al. 2006) all controlled in line with
188 previous work in the field (Taylor et al. 2011; Taylor et al. 2012a; Taylor et al. 2012b), self-reported
189 adherence was confirmed at 100% for all participants. Upon arrival, participants rested in a semi supine
190 position to provide a venous blood sample via venepuncture from the antecubital fossa for analysis of
191 eHsp72, followed by the assessment of maximal oxygen uptake [$\dot{V}O_{2max}$] as shown in Fig. 1b].
192 Participants were provided with the appropriate supplementation regime and information upon
193 departure, and were required to adhere to instructions for the following 12-weeks.

194

195 Visit 2

196

197 Following 12-weeks of supplementation, participants returned to the laboratory and followed the same
198 procedures as described for visit 1. Participants then departed the UK to undertake the MDS 2015 the
199 following week.

200

201 Visit 3

202

203 The MDS 2015 took place from 5th – 11th April, and covered a total distance of 249.4 km across the
204 Sahara Desert, Morocco (maximum temperature 39°C), over 7 days; each stage commenced at 0900h
205 (Fig. 1a indicates distances for each stage of the MDS phase of the experimental design). The MDS
206 required competitors to be self sufficient, meaning they were to carry their own food (minimum 2000

207 kcal per day), equipment, and sleeping materials for the duration of the race. Water was rationed to ~
208 9.0 – 10.5 L/day per competitor, dependent on the distance of the stage.

209

210 Upon completion of the race, participants boarded coaches and were taken from the Desert back to the
211 city of Ouarzazate, for post-race data collection (Fig. 1c). A team of trained experimenters collected
212 post-race venous blood samples; this data collection took place 6 - 8 h post-race completion.

213

214 Visit 4

215

216 Finally, participants attended the sport science laboratories 7 days post-race, whereby venous blood
217 samples and body composition were measured in line with visits 1 and 2. Participants were not
218 required to undertake a $\dot{V}O_{2\max}$ test during the final visit.

219

220 ****Please insert Fig. 1a-c here****

221

222 $\dot{V}O_{2\max}$ Test

223

224 A graded exercise test to maximal exhaustion on a motorized treadmill (Pulsar, HP Cosmos, UK)
225 began with a 5 min self-paced warm up. Thereafter, speed was increased by 1 km/h⁻¹ every 2 min, after
226 4 stages (8 min) speed remained constant and treadmill incline increased by 1% every 2 min until
227 volitional exhaustion (Winter et al. 2006). Online breath-by-breath analysis (Metalyser 3B, Cortex,
228 UK) was used to determine $\dot{V}O_{2\max}$. Measures of HR (Polar, FS1, UK) and ratings of perceived
229 exertion (RPE) were recorded every 2 min to be used as secondary criteria. The $\dot{V}O_{2\max}$ was
230 considered as the highest $\dot{V}O_2$ obtained in any 10 s period, and in line with end point criteria guidelines
231 of the ACSM; which required participants to meet a plateau in $\dot{V}O_2$, plus 2 of the 3 following criteria: a
232 failure of HR to increase with increasing exercise intensity, respiratory exchange ratio (RER) of > 1.15,
233 and RPE > 17 (ACSM 2013).

234

235 Blood collection and analysis

236

237 Venous blood samples were collected at all four visits from the antecubital fossa via venepuncture
238 (Safety blood collection set and holder, Vacuette®, Greiner Bio-One, UK), directly into three separate
239 Vacuette® tubes (4 ml; Vacuette® Greiner Bio-One, UK) treated with K3 Ethylenediaminetetraacetic
240 acid (EDTA) coagulant. Whole blood samples were centrifuged (EBA 200, Hettich, Germany) at 3000
241 rpm for 10 min for plasma separation, after which the plasma was aliquoted and stored at - 80°C until
242 analysed in duplicate for eHsp72 using a commercially available high-sensitivity enzyme-linked
243 immunosorbent assay (ELISA) kit. Whilst the measurement of plasma volume change would have been
244 beneficial, the applied nature of the study and the logistical challenges present precluded the analysis of
245 this variable. However, previous research has demonstrated that long-endurance exercise causes no
246 significant increase in haemoglobin and haematocrit values (Gomez-Merino et al. 2006; Ruell et al.
247 2006; Suzuki et al. 2006), thus the changes may have been negligible (Suzuki et al. 2000; Suzuki et al.
248 2003). Incubation of the 96-well kit was performed on an incubital shaker (Heidolph Titramax 1000,
249 Fischer Scientific, UK) at 500 rpm, and read by a plate reader (VICTOR™ X, Perkin Elmer, UK) using
250 absorption at 450 nm. Intra-assay variability was 9.93% which was in line with previous work in the
251 field: Campisi et al. (2003): < 10%; Gibson et al. (2014): 10.5%; Périard et al. (2012): 5%; Walsh et
252 al. (2001): < 10%; Whitham et al. (2006): 6.3%.

253

254 Statistical analysis

255

256 All statistical analyses were performed using the ‘psych’, ‘nlme’ and ‘stats’ packages in R version 3.3.2
257 (R Core Development Team, 2014). Normality assumptions were checked using quantile-quantile
258 plots. Normally distributed data is presented as mean standard deviation (SD), and non-normally
259 distributed data is presented as median and range. The Akaike information criteria (AIC) was used to
260 determine fit of the full model relative to the null model (Akaike 1976). A linear mixed model with
261 fixed (‘condition’, ‘time’) and random (‘subject id’) effects was fitted with a compound symmetric
262 correlation structure to determine the effect of PRO and PGLn on eHsp72 compared with CON [Time
263 (4 levels): baseline, pre-race, 6-8 h post-race, 7-d post-race × Condition (3 levels): CON, PRO, PGLn].
264 In accordance with previous literature, eHsp72 concentrations were also presented as a percentage
265 change from baseline, to account for high individual variance in baseline values and responses (Suzuki
266 et al. 2006; Morton et al. 2007; Sandström et al. 2008; Taylor et al. 2010a; Taylor et al. 2010b; Peart et

al. 2011; Gibson et al. 2014). The sex of the participants (male or female) was not included as an independent variable due to the low number of females in the experiment (4 females, 30 males). A one-way analysis of variance (ANOVA) was used to compare time to completion (minutes) between the CON, PRO, and PGLn groups. The two-tailed alpha level of significance was set as $p \leq 0.05$, and 95% confidence intervals are presented to denote the imprecision in the point estimate.

Results

Anthropometric data

Participant characteristics are displayed in Table 1. For height, those in the PRO group were 5.3 ± 1.5 cm and 6.5 ± 1.6 cm taller than those in the CON and PGLn group, respectively ($F_{2, 125} = 9.372$, $p < 0.001$). For body mass at baseline, the mass of those in the PGLn group was 8.17 ± 2.46 kg and 8.50 ± 2.57 kg less than those in the PRO and CON group, respectively ($F_{2, 125} = 7.196$, $p = 0.001$). For body mass at pre-race, the mass of those in the PGLn group was 7.20 ± 2.35 kg and 7.40 ± 2.45 kg less than those in the PRO and CON group, respectively ($F_{2, 125} = 5.992$, $p = 0.003$). For $\dot{V}O_{2\max}$ at baseline, those in the PGLn group had a score 5.33 ± 2.00 ml.kg.min⁻¹ greater than those in the CON group ($F_{2, 125} = 3.909$, $p = 0.023$). There were no differences between groups for age and $\dot{V}O_{2\max}$ at pre-race ($p > 0.05$).

eHsp72

The eHsp72 response was not different between nutritional groups (CON, PRO, PGLn). There was no main effect for group ($F_{1,2} = 3.252$, $p = 0.053$) or interaction (condition $\times$ time) effect ($F_{1,6} = 1.051$, $p = 0.399$) for eHsp72 responses. There was a main effect for time ($F_{1,3} = 22.716$, $p < 0.001$), showing that eHsp72 was elevated by 0.81 ± 0.36 ng·ml⁻¹ (95% CI = 0.20 to 1.43 ng·ml⁻¹) at 6-8 h post-race compared with baseline ($p = 0.02$). The boxplots in Fig. 2 show the variability in the responses for each group over time. Table 2 displays the mean $\pm$ SD (range) of eHsp72 for each condition at each time point. Table 3 displays the model's fixed effects coefficients and random effect variances. Individual

296 responses over time are displayed in Fig. 3 a-c, where the change is expressed as a % change from
297 baseline.

298

299 Time to completion

300

301 The time to completion was not different ($F_{1,2} = 0.615, p = 0.548$) between the CON (2819 ± 846 min,
302 95% CI = 2213 to 3425 min), PRO (2481 ± 525 min, 95% CI = 2105 to 3424 min), and PGLn ($2570 \pm$
303 707 min, 95% CI = 2064 to 3075 min) groups. The boxplots in Fig. 4 show the variability in time to
304 completion for each group.

305

306 **** Insert Table 2 near here please ****

307 ****Insert Table 3 near here please****

308

309 **** Insert Fig 2, 3 a-c and 4 near here please ****

310

311 **Discussion**

312

313 The primary aim of this study was to investigate the effects of chronic PRO and PGLn commercially
314 available supplementation on the eHsp72 response to the MDS. The experimental hypothesis was not
315 accepted in this paper, as PRO and PGLn supplementation had no effect on eHsp72 concentration over
316 the 12-week supplementation period. In addition, post-MDS eHsp72 concentration was significantly
317 elevated in all groups, however there was no difference in the magnitude of change in the CON group
318 compared to PRO or PGLn. These findings indicate that 12-week supplementation with PRO or PGLn
319 had no influence on eHsp72 concentrations.

320

321 A mean 124% increase in eHsp72 concentration was seen across all conditions post-MDS, compared to
322 pre-MDS values (Fig. 2a), supporting previous data related to the EHS response (Walsh et al. 2001;
323 Gomez-Merino et al. 2006; Suzuki et al. 2006; Gibson et al. 2014). The magnitude of this response
324 post-MDS was not different between groups, indicating that the utilised nutritional intervention had no
325 influence on pre-MDS eHsp72 concentrations, despite a likely increase in iHsp72 pre-race (Tao et al.

2006). The novelty of the present study (i.e. MDS exercise model and 12-week nutritional interventions) renders it challenging to make direct comparisons to previous research. Indeed, to the author's knowledge only two previous studies have reported the eHsp72 response to ultra-endurance exercise performance (Gomez-Merino et al. 2006; Suzuki et al. 2006). These studies demonstrated an increased eHsp72 concentration of ~ 2200% (Suzuki et al. 2006) and ~ 1674% (Gomez-Merino et al. 2006) following an ironman triathlon and 100 km run, respectively.

The difference in magnitude of response between the present study (~124% increase) and others (Gomez-Merino et al. 2006; Suzuki et al. 2006) could be attributed to a number of factors. Firstly, the logistical issues associated with field-based research, particularly the MDS location, determined that post-exercise sampling was only feasible 6-8 h post-MDS. It is therefore possible that the greatest phase of the response was not recorded. Fehrenbach et al. (2005) demonstrated that eHsp72 concentration returned to baseline levels within 24 h of exercise completion, following a marathon run, with values already significantly different at 3 h post-exercise in comparison to immediately post-exercise. It is therefore likely that the 6-8 h time course between MDS completion and sample collection in the present study elicited lower eHsp72 concentration changes from pre-race in comparison to the values that would have been displayed immediately post-race. Future research should aim to collect data as close to race completion as possible to obtain the most representative results.

Secondly, the difference may be due to a habituation effect, given the consecutive and multi-day nature of the MDS compared to the discrete within one-day ironman triathlon and 100 km run. Following 7 consecutive days of EHS, participants would likely begin to undergo heat acclimatisation, given heat acclimation/acclimatisation has been shown to commence in as few as 4 exercise-heat exposures (Petersen et al. 2010), with 'full' heat acclimation taking typically 7-14 days (Tyler et al. 2016). Thus, by the 7th consecutive day of EHS, it is likely that participants would have developed a level of acclimation to these conditions. Consequently, resting eHsp72 concentration prior to the commencement of the final MDS stage would potentially have been greater than recorded pre-race values. For example, Sandström et al. (2008) investigated the effects of 15 consecutive days heat acclimation on eHsp70, and demonstrated the impact of acclimation upon eHsp70 concentration. In

that paper, resting eHsp70 following 7 days heat acclimation was 45% greater than baseline concentration. In addition, the pre- to post-acclimation change on day 1 (54%) showed a greater effect in comparison to that of day 7 (-4.5%). This speculation could aid understanding regarding the less substantial increase in eHsp72 post-exercise in comparison to previous ultra-endurance research (Gomez-Merino et al. 2006; Suzuki et al. 2006). It is possible that the MDS induced a level of acclimatisation, which subsequently impacted the eHsp72 concentrations recorded following completion of the race, masking any influence of the probiotic supplement. However, as all participants completed the race, it is likely that the possible heat acclimatisation stimulus of MDS completion, and any subsequent confounding effects of such acclimatisation, would have been similar across participants. Whether within-race heat acclimatisation did mask any potential benefits of the probiotics cannot be excluded within the present design. Future research is recommended to analyse additional markers (such as IL-6, TNF- α), in order to obtain a greater understanding regarding the response to ultra-endurance exercise and probiotic supplementation.

Previous *in vitro* research has suggested that a key mechanism behind the protective role of probiotic (Petrof et al. 2004; Tao et al. 2006) and glutamine (Wischmeyer et al. 2001) is elevations in basal iHsp72 concentrations. Tao et al. (2006) stated that the induction of iHsp72 following probiotic treatment *in vitro* may be due to changes in gene transcription through binding of heat shock factor-1, a key activator responsible for transcription of heat shock genes (Zuhl et al. 2014). Because the systematic concentrations of Hsp72 may be proportional to the iHsp72 concentration, it was important to investigate if probiotic supplementation also influenced eHsp72 concentrations. In contrast to previous *in vitro* research (Petrof et al. 2004; Tao et al. 2006), the present study utilised commercially available supplements, which may have contained additional ingredients and subsequently impacted the response (Maughan 2005). Future research utilising similar experimental designs should develop a standardised and pharmacologically optimised probiotic supplement strain (which may likely be hybridised), to avoid the previously documented lack of quality control of probiotics (Tuomola et al. 2001). It may also be suggested that the prescribed glutamine dosage was insufficient to induce a significant eHsp72 response *in vivo*. Previous research has utilised a dosage relative to the individuals' body mass (Zuhl et al. 2014; Zuhl et al. 2015), however the present study implemented the suppliers' recommended dosage, and subsequently may have been insufficient to induce a significant response.

Should the present study have implemented the individualised dosages as utilised by Zuhl et al. (2014), dosages would range from 19.75 – 67.05 g PRO/PGLn supplementation per day (based on fat free mass range 21.95 – 74.5 kg of participants in the present study). However, only 0.9 g/day was administered in this study, demonstrating a deviation of 18.85 – 66.15 g/day from the aforementioned dosage (Zuhl et al. 2014). Evidently, these differences are substantial, and thus the effect of PRO and PGLn from the dosage utilised in the present study may have been insufficient to induce a significant response. It is important to note, however, that competitors would have utilised the commercially available dietary supplements in the same manner (i.e. in line with the supplier recommendations) as the design employed by the present study. Another consideration is the makeup of the probiotic supplement. Whilst there are limited studies in this area (particularly using the specific strains supplied in the present study), the predominance of *Lactobacillus* and *Bifidobacterium bifidum* species in this supplement indicate that these would have the strongest influence upon the eHsp72 response (if any). Although the *Lactobacillus* and *Bifidobacterium* strains activate the innate immune system (Bellavia et al. 2013; Lescheid 2014), it is unclear what specific strains interact with eHsp72. Additionally, whilst there was a level of prebiotic included within the supplement, this very small dosage (55.8 mg.d-1 fructooligosaccharides; FOS) is unlikely to have had an effect on eHsp72 concentration.

It is evident that participation in a multi-day ultra-endurance event imposes a significant level of stress upon the human body, leading to an increase in eHsp72 concentration. Such increases are likely for stimulation of pro-inflammatory cytokine and chemokines, which is typical during EHS. However, the implementation of commercially available PRO and PGLn supplementation, in line with supplier recommendations, did not increase basal eHsp72 concentration prior to the race, or alter the eHsp72 response seen in all groups post-race.

Experimental limitations, practical applications and future research directions

Collection of data within an applied setting involves a number of challenges, particularly when in a harsh environment, such as the Sahara Desert. Due to methodological limitations in the present work, it is recommended that future research collect samples immediately / within 1 h of race completion, and should implement a dosage relative to the body mass of the individual taking part, in addition to

utilising robustly standardised and optimised nutritional supplements. This will confirm if PRO or PGLn increases basal eHsp72 values prior to ultra-endurance events, particularly those completed within thermally challenging environments. Additionally, future research should focus on controlling the outlined confounding variables alongside analysis of additional biological markers, such as inflammatory markers (IL-6, TNF- α), intracellular markers (iHsp72), and markers of endotoxemia/gut damage (LPS); to greater understand the implicated biological cascades such as gut damage. To increase internal validity within similar research designs, all participants should complete the exact same robust heat acclimation regime (or completely avoid any acclimation) prior to leaving for the MDS, to control the potential confounding influence of heat acclimation/acclimatisation upon eHsp72 concentrations pre and during race. Therefore, whilst the presented results should be interpreted carefully relative to these limitations, the findings remain relevant to athletes and coaches, given we have shown that the specific commercially available probiotic supplements and chronic administration strategies utilised are unlikely to interfere with the pathways regulating immune responses to multi-day ultra-endurance running exercise in the heat.

430

431 **Acknowledgements**

432

433 The authors would like to thank BioCare UK for providing the nutritional supplements used in the
434 study, RunUltra for allowing access to the competitors, and the participants for their cooperation.

435

436 **References**

437

438 ACSM (2013). ACSM's guidelines for exercise testing and prescription: Lippincott Williams &
439 Wilkins.

440 Adams, W.M., Hosokawa, Y., and Casa, D.J. (2015). The Timing of Exertional Heat Stroke Survival
441 Starts prior to Collapse. *Current Sports Medicine Reports* **14**: 273-274.
442 doi:10.1249/jsr.000000000000166.

443 Akaike, H. (1976). Canonical Correlation Analysis of Time Series and the Use of an Information
444 Criterion. *In: Mathematics in Science and Engineering. Edited by K.M. Raman and G.L. Dimitri*, pp.
445 27-96: Elsevier.

- 446 Amorim, F.T., Yamada, P.M., Robergs, R.A., Schneider, S.M., and Moseley, P.L. (2008). The effect of
447 the rate of heat storage on serum heat shock protein 72 in humans. *European Journal Of Applied*
448 *Physiology* **104**: 965-972. doi:10.1007/s00421-008-0850-3.
- 449 Anbarasi, K., Kathirvel, G., Vani, G., Jayaraman, G., and Devi, C.S. (2006). Cigarette smoking induces
450 heat shock protein 70 kDa expression and apoptosis in rat brain: modulation by baclofen A.
451 *Neuroscience* **138**: 1127-1135. doi:10.1016/j.neuroscience.2005.11.029.
- 452 Armstrong, L.E., Casa, D.J., Millard-Stafford, M., Moran, D.S., Pyne, S.W., and Roberts, W.O. (2007).
453 Exertional Heat Illness during Training and Competition. *Medicine & Science in Sports & Exercise* **39**:
454 556-572. doi:10.1249/MSS.0b013e31802fa199.
- 455 Asea, A. (2005). Stress proteins and initiation of immune response: chaperone activity of hsp72.
456 *Exercise immunology review* **11**: 34.
- 457 Bellavia, M., Tomasello, G., Romeo, M., Damiani, P., Monte, A.I.L., Lozio, L., Campanella, C.,
458 Gammazza, A.M., Rappa, F., and Zummo, G. (2013). Gut microbiota imbalance and chaperoning
459 system malfunction are central to ulcerative colitis pathogenesis and can be counteracted with
460 specifically designed probiotics: a working hypothesis. *Medical microbiology and immunology* **202**:
461 393-406. doi:10.1007/s00430-013-0305-2.
- 462 Binkley, H.M., Beckett, J., Casa, D.J., Kleiner, D.M., and Plummer, P.E. (2002). National Athletic
463 Trainers' Association position statement: exertional heat illnesses. *Journal of Athletic Training* **37**: 329.
- 464 Campisi, J., Leem, T.H., and Fleshner, M. (2003). Stress-induced extracellular Hsp72 is a functionally
465 significant danger signal to the immune system. *Cell Stress & Chaperones* **8**: 272-286.
- 466 da Fonseca-Engelhardt, K., Knechtle, B., Rüst, C.A., Knechtle, P., Lepers, R., and Rosemann, T.
467 (2013). Participation and performance trends in ultra-endurance running races under extreme
468 conditions- 'Spartathlon' versus 'Badwater'. *Extreme Physiology & Medicine* **2**: 1. doi:10.1186/2046-
469 7648-2-15.
- 470 Febbraio, M.A., Steensberg, A., Walsh, R., Koukoulas, I., van Hall, G., Saltin, B., and Pedersen, B.K.
471 (2002). Reduced glycogen availability is associated with an elevation in HSP72 in contracting human
472 skeletal muscle. *Journal of Physiology* **538**: 911-917. doi:10.1013/jphysiol.2001.013145.
- 473 Fehrenbach, E., Niess, A.M., Voelker, K., Northoff, H., and Mooren, F.C. (2005). Exercise intensity
474 and duration affect blood soluble HSP72. *International Journal of Sports Medicine* **26**: 552-557.
475 doi:10.1055/s-2004-830334.

- Gibson, O.R., Dennis, A., Parfitt, T., Taylor, L., Watt, P.W., and Maxwell, N.S. (2014). Extracellular Hsp72 concentration relates to a minimum endogenous criteria during acute exercise-heat exposure. *Cell Stress Chaperones* **19**: 389-400. doi:10.1007/s12192-013-0468-1.
- Goforth, C.W. and Kazman, J.B. (2015). Exertional heat stroke in navy and marine personnel: a hot topic. *Critical care nurse* **35**: 52-59. doi:10.4037/ccn2015257.
- Gomez-Merino, D., Drogou, C., Guezennec, C.Y., Burnat, P., Bourrilhon, C., Tomaszewski, A., Milhau, S., and Chennaoui, M. (2006). Comparison of systemic cytokine responses after a long distance triathlon and a 100-km run: relationship to metabolic and inflammatory processes. *European Cytokine Network* **17**: 117-24.
- González-Alonso, J., Crandall, C.G., and Johnson, J.M. (2008). The cardiovascular challenge of exercising in the heat. *The Journal of Physiology* **586**: 45-53. doi:10.1113/jphysiol.2007.142158.
- Hillman, A.R., Vince, R.V., Taylor, L., McNaughton, L., Mitchell, N., and Siegler, J. (2011). Exercise-induced dehydration with and without environmental heat stress results in increased oxidative stress. *Appl Physiol Nutr Metab* **36**: 698-706. doi:10.1139/h11-080.
- Johnson, J.D. and Fleshner, M. (2006). Releasing signals, secretory pathways, and immune function of endogenous extracellular heat shock protein 72. *Journal of Leukocyte Biology* **79**: 425-434. doi:10.1189/jlb.0905523.
- Knechtle, B., Knechtle, P., and Lepers, R. (2011). Participation and performance trends in ultra-triathlons from 1985 to 2009. *Scandinavian Journal of Medicine & Science in Sports* **21**: e82-e90. doi:10.1111/j.1600-0838.2010.01160.x.
- Lehner, T., Bergmeier, L.A., Wang, Y., Tao, L., Sing, M., Spallek, R., and Zee, R.v.d. (2000). Heat shock proteins generate b-chemokines which function as innate adjuvants enhancing adaptive immunity. *European journal of immunology* **30**: 594-603. doi:10.1002/1521-4141(200002)30:2<594::AID-IMMU594>3.0.CO;2-1.
- Leon, L.R. and Helwig, B.G. (2010). Heat stroke: Role of the systemic inflammatory response.
- Lescheid, D.W. (2014). Probiotics as regulators of inflammation: A review. *Functional Foods in Health and Disease* **4**: 299-311.
- Lim, C. and Mackinnon, L. (2006). The Roles of Exercise-Induced Immune System Disturbances in the Pathology of Heat Stroke. *Sports Medicine* **36**: 39-64. doi:10.2165/00007256-200636010-00004.

- Lu, P.-Z., Lai, C.-Y., and Chan, W.-H. (2008). Caffeine induces cell death via activation of apoptotic signal and inactivation of survival signal in human osteoblasts. *International journal of molecular sciences* **9**: 698-718. doi:10.3390/ijms9050698.
- Magalhaes, C., Amorim, F.T., Passos, R.L., Fonseca, M.A., Oliveira, K.P., Lima, M.R., Guimaraes, J.B., Ferreira-Junior, J.B., Martini, A.R., Lima, N.R., Soares, D.D., Oliveira, E.M., and Rodrigues, L.O. (2010). Heat and exercise acclimation increases intracellular levels of Hsp72 and inhibits exercise-induced increase in intracellular and plasma Hsp72 in humans. *Cell Stress Chaperones* **15**: 885-95. doi:10.1007/s12192-010-0197-7.
- Marshall, H.C., Ferguson, R.A., and Nimmo, M.A. (2006). Human resting extracellular heat shock protein 72 concentration decreases during the initial adaptation to exercise in a hot, humid environment. *Cell Stress Chaperones* **11**: 129-34.
- Maughan, R.J. (2005). Contamination of dietary supplements and positive drug tests in sport. *Journal of Sports Sciences* **23**: 883-889. doi:10.1080/02640410400023258.
- Mizzen, L.A. and Welch, W.J. (1988). Characterization of the thermotolerant cell. I. Effects on protein synthesis activity and the regulation of heat-shock protein 70 expression. *The Journal of Cell Biology* **106**: 1105.
- Morton, J.P., MacLaren, D.P.M., Cable, N.T., Campbell, I.T., Evans, L., Bongers, T., Griffiths, R.D., Kayani, A.C., McArdle, A., and Drust, B. (2007). Elevated core and muscle temperature to levels comparable to exercise do not increase heat shock protein content of skeletal muscle of physically active men. *Acta Physiologica* **190**: 319-327. doi:10.1111/j.1748-1716.2007.01711.x.
- Nybo, L. (2008). Hyperthermia and fatigue. *Journal of Applied Physiology* **104**: 871-878. doi:10.1152/jappphysiol.00910.2007.
- Nybo, L. and Nielsen, B. (2001). Hyperthermia and central fatigue during prolonged exercise in humans.
- Nybo, L., Rasmussen, P., and Sawka, M.N. (2014). Performance in the heat-physiological factors of importance for hyperthermia-induced fatigue. *Comprehensive physiology* **4**: 657-89. doi:10.1002/cphy.c130012.
- Peart, D.J., McNaughton, L.R., Midgley, A.W., Taylor, L., Towlson, C., Madden, L.A., and Vince, R.V. (2011). Pre-exercise alkalosis attenuates the heat shock protein 72 response to a single-bout of

- anaerobic exercise. *Journal of Science and Medicine in Sport* **14**: 435-40.
doi:10.1016/j.jsams.2011.03.006.
- Périard, J.D., Ruell, P., Caillaud, C., and Thompson, M.W. (2012). Plasma Hsp72 (HSPA1A) and Hsp27 (HSPB1) expression under heat stress: influence of exercise intensity. *Cell Stress and Chaperones* **17**: 375-383. doi:10.1007/s12192-011-0313-3.
- Petersen, C.J., Portus, M.R., Pyne, D.B., Dawson, B.T., Cramer, M.N., and Kellett, A.D. (2010). Partial heat acclimation in cricketers using a 4-day high intensity cycling protocol. *Int J Sports Physiol Perform* **5**: 535-45.
- Petrof, E.O., Kojima, K., Ropeleski, M.J., Musch, M.W., Tao, Y., De Simone, C., and Chang, E.B. (2004). Probiotics inhibit nuclear factor- κ B and induce heat shock proteins in colonic epithelial cells through proteasome inhibition. *Gastroenterology* **127**: 1474-1487.
- Ruell, P.A., Thompson, M.W., Hoffman, K.M., Brotherhood, J.R., and Richards, D.A.B. (2006). Plasma Hsp72 is higher in runners with more serious symptoms of exertional heat illness. *European Journal Of Applied Physiology* **97**: 732-736.
- Sandström, M.E., Siegler, J.C., Lovell, R.J., Madden, L.A., and McNaughton, L. (2008). The effect of 15 consecutive days of heat-exercise acclimation on heat shock protein 70. *Cell Stress & Chaperones* **13**: 169-175. doi:10.1007/s12192-008-0022-8.
- Suzuki, K., Nakaji, S., Yamada, M., Liu, Q., Kurakake, S., Okamura, N., Kumae, T., Umeda, T., and Sugawara, K. (2003). Impact of a competitive marathon race on systemic cytokine and neutrophil responses. *Med Sci Sports Exerc* **35**: 348-55. doi:10.1249/01.mss.0000048861.57899.04.
- Suzuki, K., Peake, J., Nosaka, K., Okutsu, M., Abbiss, C., Surriano, R., Bishop, D., Quod, M., Lee, H., Martin, D., and Laursen, P. (2006). Changes in markers of muscle damage, inflammation and HSP70 after an Ironman triathlon race. *European Journal of Applied Physiology* **98**: 525-534. doi:10.1007/s00421-006-0296-4.
- Suzuki, K., Yamada, M., Kurakake, S., Okamura, N., Yamaya, K., Liu, Q., Kudoh, S., Kowatari, K., Nakaji, S., and Sugawara, K. (2000). Circulating cytokines and hormones with immunosuppressive but neutrophil-priming potentials rise after endurance exercise in humans. *Eur J Appl Physiol* **81**: 281-7. doi:10.1007/s004210050044.
- Tao, Y., Drabik, K.A., Waypa, T.S., Musch, M.W., Alverdy, J.C., Schneewind, O., Chang, E.B., and Petrof, E.O. (2006). Soluble factors from *Lactobacillus* GG activate MAPKs and induce cytoprotective

- 564 heat shock proteins in intestinal epithelial cells. *American Journal of Physiology-Cell Physiology* **290**:
 565 C1018-C1030. doi:10.1152/ajpcell.00131.2005.
- 566 Taylor, L., Hillman, A.R., Midgley, A.W., Peart, D.J., Christmas, B., and McNaughton, L.R. (2012a).
 567 Hypoxia-mediated prior induction of monocyte-expressed HSP72 and HSP32 provides protection to
 568 the disturbances to redox balance associated with human sub-maximal aerobic exercise. *Amino Acids*
 569 **43**: 1933-1944. doi:10.1007/s00726-012-1265-3.
- 570 Taylor, L., Midgley, A.W., Christmas, B., Hilman, A.R., Madden, L.A., Vince, R.V., and McNaughton,
 571 L.R. (2011). Daily hypoxia increases basal monocyte HSP72 expression in healthy human subjects.
 572 *Amino acids* **40**: 393-401. doi:10.1007/s00726-010-0644-x.
- 573 Taylor, L., Midgley, A.W., Christmas, B., Madden, L.A., Vince, R.V., and McNaughton, L.R. (2010a).
 574 Daily quadratic trend in basal monocyte expressed HSP72 in healthy human subjects. *Amino Acids* **38**:
 575 1483-8. doi:10.1007/s00726-009-0360-6.
- 576 Taylor, L., Midgley, A.W., Christmas, B., Madden, L.A., Vince, R.V., and McNaughton, L.R. (2010b).
 577 The effect of acute hypoxia on heat shock protein 72 expression and oxidative stress in vivo. *European*
 578 *Journal Of Applied Physiology* **109**: 849-855. doi:10.1007/s00421-010-1430-x.
- 579 Taylor, L., Midgley, A.W., Sandstrom, M.E., Christmas, B., and McNaughton, L.R. (2012b). The effect
 580 of the hyperbaric environment on heat shock protein 72 expression in vivo. *Research in sports*
 581 *medicine* **20**: 142-153. doi:10.1080/15438627.2012.660830.
- 582 Taylor, L., Watkins, S., Marshall, H., Dascombe, B., and Foster, J. (2015). The impact of different
 583 environmental conditions on cognitive function: a focused review. *Frontiers in Physiology* **6**.
 584 doi:10.3389/fphys.2015.00372.
- 585 Tuomola, E., Crittenden, R., Playne, M., Isolauri, E., and Salminen, S. (2001). Quality assurance
 586 criteria for probiotic bacteria. *The American journal of clinical nutrition* **73**: 393s-398s.
- 587 Tyler, C.J., Reeve, T., Hodges, G.J., and Cheung, S.S. (2016). The Effects of Heat Adaptation on
 588 Physiology, Perception and Exercise Performance in the Heat: A Meta-Analysis. *Sports Medicine*: 1-
 589 26. doi:10.1007/s40279-016-0538-5.
- 590 Walsh, R.C., Koukoulas, I., Garnham, A., Moseley, P.L., Hargreaves, M., and Febbraio, M.A. (2001).
 591 Exercise increases serum Hsp72 in humans. *Cell Stress Chaperones* **6**: 386-93.
- 592 Whitham, M. and Fortes, M.B. (2008). Heat shock protein 72: release and biological significance
 593 during exercise. *Frontiers in Bioscience* **13**: 1328-1339.

Whitham, M., Laing, S.J., Jackson, A., Maassen, N., and Walsh, N.P. (2007). Effect of exercise with and without a thermal clamp on the plasma heat shock protein 72 response. *Journal of Applied Physiology* **103**: 1251-1256. doi:10.1152/jappphysiol.00484.2007.

Whitham, M., Walker, G.J., and Bishop, N.C. (2006). Effect of caffeine supplementation on the extracellular heat shock protein 72 response to exercise. *Journal of Applied Physiology* **101**: 1222-1227. doi:10.1152/jappphysiol.00409.2006.

Wijerathne, B.T., Pilapitiya, S.D., Vijitharan, V., Farah, M.M., Wimalasooriya, Y.V., and Siribaddana, S.H. (2016). Exertional heat stroke in a young military trainee: is it preventable? *Military Medical Research* **3**: 1. doi:10.1186/s40779-016-0078-1.

Winter, E.M., Jones, A.M., Davison, R.R., Bromley, P.D., and Mercer, T.H. (2006). *Sport and Exercise Physiology Testing Guidelines: Volume I-Sport Testing: The British Association of Sport and Exercise Sciences Guide*: Routledge.

Wischmeyer, P.E. (2002). Glutamine and heat shock protein expression. *Nutrition* **18**: 225-228. doi:10.1016/S0899-9007(01)00796-1.

Wischmeyer, P.E., Kahana, M., Wolfson, R., Ren, H., Musch, M.M., and Chang, E.B. (2001). Glutamine induces heat shock protein and protects against endotoxin shock in the rat. *Journal of Applied Physiology* **90**: 2403-2410.

Zaryski, C. and Smith, D.J. (2005). Training principles and issues for ultra-endurance athletes. *Current Sports Medicine Reports* **4**: 165-170. doi:10.1007/s11932-005-0062-0.

Zuhl, M., Dokladny, K., Mermier, C., Schneider, S., Salgado, R., and Moseley, P. (2015). The effects of acute oral glutamine supplementation on exercise-induced gastrointestinal permeability and heat shock protein expression in peripheral blood mononuclear cells. *Cell Stress & Chaperones* **20**: 85-93. doi:10.1007/s12192-014-0528-1.

Zuhl, M., Lanphere, K., Kravitz, L., Mermier, C., Schneider, S., Dokladny, K., and Moseley, P. (2014). Effects of oral glutamine supplementation on exercise-induced gastrointestinal permeability and tight junction protein expression. *Journal of Applied Physiology* **116**: 183-191. doi:10.1152/jappphysiol.00646.2013.

621
622
623
624
625
626

Table 1. Anthropometrical characteristics of age, height, mass; body fat percentage, and $\dot{V}O_{2\max}$ data (week 1 and week 12) are reported as means $\pm$ SD (age is reported as range due to non-normal distribution).

Group	Age (years)	Baseline		Pre-race	
		Mass (Kg)	$\dot{V}O_{2\max}$ (ml.kg.min ⁻¹)	Mass (Kg)	$\dot{V}O_{2\max}$ (ml.kg.min ⁻¹)
PRO	25 – 50	78.87 $\pm$ 6.93	56.42 $\pm$ 8.26	77.49 $\pm$ 6.52	59.40 $\pm$ 6.33
PGLn	31 – 53	71.34 $\pm$ 12.92	56.50 $\pm$ 5.95	70.09 $\pm$ 12.31	59.25 $\pm$ 6.92
CON	23 – 60	79.50 $\pm$ 14.07	51.17 $\pm$ 12.68	77.29 $\pm$ 12.92	55.25 $\pm$ 11.96

Table 2. Mean $\pm$ SD (range) eHsp72 (ng.ml⁻¹) concentration between groups at each data collection time point.

Group	Baseline	Pre-race	6 hours post race	7 d post-race
CON	1.68 $\pm$ 1.65 (0.59 – 3.81)	1.82 $\pm$ 1.98 (0.63 – 1.93)	3.45 $\pm$ 3.35 (0.78 – 3.30)	2.05 $\pm$ 2.49 (0.67 – 2.17)
PRO	1.20 $\pm$ 0.58 (0.68 – 2.70)	1.58 $\pm$ 0.87 (0.75 – 2.96)	2.77 $\pm$ 1.21 (1.82 – 7.30)	1.80 $\pm$ 1.27 (0.82 – 4.13)
PGLn	1.77 $\pm$ 2.51 (0.59 – 1.90)	1.03 $\pm$ 0.45 (0.73 – 2.19)	2.20 $\pm$ 0.76 (1.57 – 4.16)	1.04 $\pm$ 0.41 (0.71 – 2.16)

Kg: kilograms; **$\dot{V}O_{2\max}$:** maximal volume of oxygen consumption; **ml.kg.min⁻¹:** millilitres per kilogram per minute. **PRO:** Probiotic; **PGLn:** Probiotic + Glutamine; **CON:** Control

Table 3. Beta coefficients (*B*), 95 % confidence intervals (*CI*), and alpha values (*p*) are reported for the fixed components (Supplement & time) before and after the MDS race. The standard deviation of the intercept and residual are reported for the random effect (subject ID).

	eHsp72 (ng.ml ⁻¹)		
	<i>B</i>	95% <i>CI</i>	<i>p</i>
Fixed Parts			
(Intercept)	1.31	0.79 – 1.83	<.001
Condition Effect			
PRO	0.02	-0.72 – 0.75	0.968
PGLn	-0.32	-1.09 – 0.45	0.421
Time Effect			
Pre-Race	-0.1	-0.70 – 0.50	0.742
6-8 h Post-Race	0.81	0.20 – 1.43	0.011
1-Week Post-Race	-0.04	-0.67 – 0.60	0.911
Condition: Time Interaction			
PRO: Pre-Race	0.27	-0.59 – 1.13	0.541
PGLn: Pre-Race	0.13	-0.75 – 1.01	0.770
PRO: 6-8 h Post-Race	1.01	0.15 – 1.87	0.023
PGLn: 6-8 h Post-Race	0.38	-0.51 – 1.28	0.401
PRO: 1-Week Post-Race	0.6	-0.31 – 1.50	0.198
PGLn: 1-Week Post-Race	0.09	-0.82 – 1.00	0.851
Random Parts			
σ^2		0.511	
$\tau_{00, ID}$		0.258	

673 **Figure legends**

674

675 **Fig. 1 a-c** Experimental schematic outlining: a: the full experimental study design, b: procedures of
676 Visits 1, 2 and 4, c: procedure of Visit 3

677

678 **Fig. 2** Boxplots showing the variability in responses for each group over time. #Denotes a significant
679 effect of time on eHsp72 concentration

680

681 **Fig. 3 a-c** Individual eHsp72 responses within groups, as a % change from baseline (a: PRO, b: PGLn,
682 c: CON)

683

684 **Fig. 4** Boxplots showing the variability in time to completion for each group

Draft

Fig. 1 a-c Experimental schematic outlining: a: the full experimental study design, b: procedures of Visits 1, 2 and 4, c: procedure of Visit 3

254x190mm (200 x 200 DPI)

Fig. 2 Boxplots showing the variability in responses for each group over time. #Denotes a significant effect of time on eHsp72 concentration

336x222mm (96 x 96 DPI)

Fig. 3 a-c Individual eHsp72 responses within groups, as a % change from baseline (a: PRO, b: PGLn, c: CON)

110x233mm (300 x 300 DPI)

Fig. 4 Boxplots showing the variability in time to completion for each group

136x116mm (96 x 96 DPI)