



## Repression of Hedgehog signal transduction in T-lineage cells increases TCR-induced activation and proliferation

Nicola J. Rowbotham, Anna L. Furmanski, Ariadne L. Hager-Theodorides, Susan E. Ross, Ekati Drakopoulou, Costas Koufaris, Susan V. Outram & Tessa Crompton

To cite this article: Nicola J. Rowbotham, Anna L. Furmanski, Ariadne L. Hager-Theodorides, Susan E. Ross, Ekati Drakopoulou, Costas Koufaris, Susan V. Outram & Tessa Crompton (2008) Repression of Hedgehog signal transduction in T-lineage cells increases TCR-induced activation and proliferation, *Cell Cycle*, 7:7, 904-908, DOI: [10.4161/cc.7.7.5628](https://doi.org/10.4161/cc.7.7.5628)

To link to this article: <https://doi.org/10.4161/cc.7.7.5628>



Copyright © 2008 Landes Bioscience



Published online: 16 Apr 2008.



Submit your article to this journal [↗](#)



Article views: 120



View related articles [↗](#)



Citing articles: 15 View citing articles [↗](#)

## Brief Report

# Repression of hedgehog signal transduction in T-lineage cells increases TCR-induced activation and proliferation

Nicola J. Rowbotham,<sup>1,2</sup> Anna L. Furmanski,<sup>1</sup> Ariadne L. Hager-Theodorides,<sup>1</sup> Susan E. Ross,<sup>1,2</sup> Ekati Drakopoulou,<sup>1</sup> Costas Koufaris,<sup>2</sup> Susan V. Outram<sup>1</sup> and Tessa Crompton<sup>1,2</sup>

<sup>1</sup>Immunobiology Unit; UCL Institute of Child Health; London, United Kingdom; <sup>2</sup>Division of Cell and Molecular Biology; Imperial College London; London, United Kingdom

**Key words:** hedgehog signalling, Gli2, Shh, T cell, thymocyte development, T cell activation, proliferation

Hedgehog proteins signal for differentiation, survival and proliferation of the earliest thymocyte progenitors, but their functions at later stages of thymocyte development and in peripheral T-cell function are controversial. Here we show that repression of Hedgehog (Hh) pathway activation in T-lineage cells, by expression of a transgenic repressor form of Gli2 (Gli2 $\Delta$ C2), increased T-cell differentiation and activation in response to TCR signalling. Expression of the Gli2 $\Delta$ C2 transgene increased differentiation from CD4<sup>+</sup>CD8<sup>+</sup> to single positive thymocyte, and increased peripheral T cell populations. Gli2 $\Delta$ C2 T-cells were hyper-responsive to activation by ligation of CD3 and CD28: they expressed cell surface activation markers CD69 and CD25 more quickly, and proliferated more than wild-type T-cells. These data show that Hedgehog pathway activation in thymocytes and T-cells negatively regulates TCR-dependent differentiation and proliferation. Thus, as negative regulators of TCR-dependent events, Hh proteins provide an environmental influence on T-cell fate.

## Introduction

The Hedgehog (Hh) family of secreted intercellular signalling molecules regulate both the embryonic development and adult homeostasis of many mammalian tissues, including the immune system.<sup>1-4</sup> Hh proteins bind to their cells surface receptor Patched (Ptch) releasing the signal transducer Smoothened (Smo) to transmit the Hh signal into the cell, leading to transcriptional changes mediated by transcription factors Gli1, Gli2 and Gli3.<sup>2</sup> Gli1 acts exclusively as an activator of transcription, and is neither essential for mouse development nor initiation of the Hh signal.<sup>5</sup> Gli2 and Gli3 are both essential for mouse development and can undergo processing to act as transcriptional activators in the presence of Hh, or repressors in its absence.<sup>6,7</sup> Gli2 is necessary to initiate the first transcriptional changes upon Hh signalling.<sup>8</sup>

In the thymus, analysis of mouse mutants of Sonic Hh (Shh), Gli3 and Smo have demonstrated an essential positive regulatory

role for the pathway in signalling for survival, proliferation and differentiation of the earliest thymocyte progenitor populations.<sup>9-11</sup> The influence of the pathway at later stages of T lineage differentiation and on T cell activation, however, is highly controversial, with different experimental models suggesting variously opposing effects, or no effect.<sup>9,12-16</sup> In vitro addition of recombinant Shh (r-Shh) to T cells has been shown to enhance T cell activation and proliferation induced by antibodies against CD3 and CD28.<sup>13,14</sup> In contrast, constitutive activation of Hh signalling by expression of a transgenic activator form of Gli2 (Gli2 $\Delta$ N<sub>2</sub>) in T-lineage cells inhibited T cell activation and proliferation, by repressing TCR signal transduction.<sup>3,12</sup> Surprisingly however, a recent study in which Smo was conditionally deleted from T-lineage cells failed to reveal any influence (either positive or negative) of loss of Hh signalling on anti-CD3 induced T cell proliferation.<sup>9</sup>

Here we describe a novel mouse model in which Hh signalling is repressed in T-lineage cells, by transgenic expression of the repressor form of Gli2 (Gli2 $\Delta$ C2), in order to ask if T cell autonomous inhibition of Hh signalling influences the TCR dependent stages of T cell differentiation and peripheral T cell function.

## Results

**Transgenic Gli2 $\Delta$ C<sub>2</sub> expression in thymocytes leads to a repression in Hh signalling.** To test the effect of T cell autonomous inhibition of Hh signalling on T cell development and T cell activation, we produced a transgenic mouse that expressed a C-terminally truncated form of Gli2 under the control of the Ick promoter. Transgene expression was thus restricted to T-lineage cells from the DN2 stage onwards (with full expression achieved at DN4), including peripheral T cells, but not thymic epithelium.<sup>17</sup> This truncated form of Gli2 (Gli2 $\Delta$ C<sub>2</sub>) can only act as a strong repressor of transcription, thus mimicking the transcriptional events that occur in the absence of Hh signalling.<sup>6</sup> Two independent transgenic lines were generated with similar copy number (Fig. 1A). Both lines showed the same phenotype so we present experimental data from one line only. Transgene expression was demonstrated by quantitative RT-PCR in both thymus and periphery, with expression levels corresponding to the proportion of T-lineage cells (Fig. 1B). To confirm that the Gli2 $\Delta$ C<sub>2</sub> transgene was functional and able to repress an exogenous Hh signal, we compared the ability of purified CD4<sup>+</sup> T cells to upregulate the Hh target gene *Ptc1* in response to

\*Correspondence to: Tessa Crompton; Immunobiology Unit; UCL Institute of Child Health; 30 Guilford Street; London WC1N 1EH UK; Tel.: +44.(0)207.905.2893; Fax: +44.(0)207.813.8494; Email: t.crompton@ich.ucl.ac.uk

Submitted: 01/16/08; Accepted: 01/18/08

Previously published online as a *Cell Cycle* E-publication:  
<http://www.landesbioscience.com/journals/cc/article/5628>




Figure 1. For figure legend, see page 906.

Figure 1. Phenotype of Gli2 $\Delta$ C<sub>2</sub> mice. (A) Quantitative RT-PCR to show transgene copy number for both transgene positive lines. Error bars show standard deviation (SD). (B) Gli2 expression relative to HPRT expression by quantitative RT-PCR analysis in thymus and spleen of Gli2 $\Delta$ C<sub>2</sub> transgenic tissues. Error bars show SD. (C) Ptc1 expression relative to HPRT expression as assessed by quantitative RT-PCR analysis in purified CD4<sup>+</sup> Gli2 $\Delta$ C<sub>2</sub> or WT splenocytes cultured with or without 0.2  $\mu$ g r-Shh for 3 hours. Gli2 $\Delta$ C<sub>2</sub> cells are less responsive to Shh signalling than WT cells. Error bars show SD. (D) Relative cell number (calculated relative to the mean of WT littermates) in thymus and spleen. (E) Histograms to show CD3 expression in thymus and peripheral CD4 and CD8 SP cells. Numbers indicate percentage of cells falling within the marker. (F) Typical CD4 and CD8 profile of thymus (top panel) and spleen (lower panel) of a Gli2 $\Delta$ C<sub>2</sub> transgenic mouse and its littermate. Numbers indicate the percentage of cells falling within each quadrant. Bar charts show the relative proportion of different populations. Differences were statistically significant as given: (thymus) CD4SP,  $p = .049$ ; CD8SP,  $p = .005$ ; DP,  $p = .017$ ; and (spleen) CD4SP,  $p = .024$ ; CD8SP,  $p = .007$ .

treatment with r-Shh. There was a marked decrease in the ability of the Gli2 $\Delta$ C<sub>2</sub> transgenic CD4<sup>+</sup> cells to upregulate Ptc1 confirming that the Gli2 $\Delta$ C<sub>2</sub> protein was a successful repressor of Hh signalling (Fig. 1C).

**Inhibition of Hh signalling in the Gli2 $\Delta$ C<sub>2</sub> transgenic influences the production of SP cells.** Initial analysis showed that thymus and spleen were grossly normal, with no marked differences in lymphocyte numbers or TCR/CD3 expression in the thymus or spleen between WT and Gli2 $\Delta$ C<sub>2</sub> (Fig. 1D and E). In the thymus there was, however, a significant increase in the proportion of SP cells, with concomitant decrease in the DP population, indicating that reduction in the ability of the thymocyte to respond to the Hh signal increases differentiation from DP to SP cell (Fig. 1E).

The effect of reduction of Hh signal at the transition from DP to SP thymocyte is controversial. In the Shh<sup>-/-</sup> thymus there is an increase in the production of SP cells.<sup>12</sup> Consistent with this, T-lineage specific constitutive activation of Hh signalling decreased DP to SP transition.<sup>12</sup> In contrast, in the T-lineage specific conditional Smo knock-out, no effect on this transition was described,<sup>9</sup> suggesting that the action of Shh on the production of SP cells could be an indirect effect through another cell type. However, the repression of Hh signalling described here is restricted to thymocytes through use of the lck transgenic cassette, demonstrating a T cell autonomous effect of Hh signalling on DP to SP differentiation.

In peripheral lymphoid organs we observed a significant increase in T cell populations (Fig. 1E).

**Increased T cell activation in peripheral Gli2 $\Delta$ C<sub>2</sub> transgenic T cells.** To test the effect of repression of Hh signalling on T cell activation, splenocytes were activated with anti-CD3 and anti-CD28. After 24 hours we observed an increase in expression of the early activation marker CD69 and the later activation marker CD25, in both CD4<sup>+</sup> and CD8<sup>+</sup> cells from the Gli2 $\Delta$ C<sub>2</sub> transgenic compared to WT littermates (Fig. 2A and B).

**Enhanced proliferation of Gli2 $\Delta$ C<sub>2</sub> T Cells on TCR/CD28 ligation.** To assess proliferation, WT and Gli2 $\Delta$ C<sub>2</sub> transgenic splenocytes were labelled with CFSE and stimulated with anti-CD3 and anti-CD28. The Gli2 $\Delta$ C<sub>2</sub> T cells underwent more cell divisions than their WT counterparts (Fig. 2C) demonstrating that repression of full Hh pathway activation in T cells increases their ability to proliferate in response to TCR/CD28 ligation.

To ask if the increase in proliferation was saturated at this level of Gli2 $\Delta$ C<sub>2</sub> expression, we doubled the transgene copy number by crossing the transgenic mice with each other. This further increased proliferation (Fig. 2D). The fact that we were able to increase repression demonstrates competitive inhibition of Hh-dependent transcription by the Gli2 $\Delta$ C<sub>2</sub> transgene.

## Discussion

Loss-of-function mutants have previously shown that Hh signalling promotes proliferation of very early thymocytes progenitors.<sup>9,11</sup> In contrast, here we show that constitutive repression of the Hh pathway promotes proliferation of mature T cells. The effect of Hh signalling on proliferation has also appeared ambiguous in other systems.<sup>18</sup> In the chick and mouse retina Hh pathway activation has been shown to promote proliferation,<sup>19-21</sup> whereas Hh mutant Zebrafish have prolonged retinal proliferation, due to the inability of precursor cells to exit the cell cycle.<sup>22</sup> Recent work has suggested that Hh signalling may effect stem, progenitor and mature cells differently, with Hh signalling either promoting cell cycle progression or pushing cells out of the cell cycle, depending on the state of differentiation of the cell.<sup>23</sup> The T cell lineage thus provides another example of Hh activation promoting proliferation of the progenitor cell, but limiting the proliferation of the more differentiated mature cell.

In summary, here we have shown that repression of Hh pathway activation in thymocytes and T cells increases differentiation and activation in response to TCR signalling. Thus, Hh proteins are negative regulators of TCR-dependent events in T cells, providing an environmental influence on T cell fate.

## Methods

**Construction of lck-Gli2 $\Delta$ N<sub>2</sub> transgenic.** Gli2 $\Delta$ C<sub>2</sub> cDNA<sup>6</sup> was bluntend cloned into the BamHI site of the lck proximal promoter cassette.<sup>24</sup> The 8.9 kb transgene was isolated by *NotI* digestion and purified using QIAEXII Gel Extraction Kit (Qiagen). CBA x C57BL/6 oocytes were injected, generating 2 independent transgenic lines.

**Mice.** lck-Gli2 $\Delta$ C<sub>2</sub> transgene-positive mice were backcrossed for >7 generations with C57BL/6 (B&K Universal) and maintained under UK Home Office regulations.

**Genotyping.** DNA was extracted as described.<sup>11</sup> Gli2 $\Delta$ C<sub>2</sub> transgene positive mice were detected by the presence of human growth hormone DNA (5'-hGH CGAACCCTCAGGGTCCTGTGG, 3'-hGH GGATTTCTGTTGTGTTTCCTCCCTG).<sup>25</sup>

**EasySep bead purification.** CD4<sup>+</sup> lymphocytes were purified from whole spleen by magnetic bead separation using the EasySep<sup>®</sup> Negative Selection Mouse CD4<sup>+</sup> T cell Enrichment kit (StemCell Technologies, UK) according to the manufacturer's instructions giving  $\geq 95\%$  purity.

**Quantitative RT-PCR.** Quantitative RT-PCR was as described.<sup>10</sup> Primers were Gli2 $\Delta$ C<sub>2</sub>F:AGAACCCTGAAGACACACCTGCG, Gli2 $\Delta$ C<sub>2</sub>R:GAGGCATTGGAGAAGGCTTTG. Ptc1F:TGCTCTCCAGTTCTCAGACTC, Ptc1R:CCACAACCTTGGCTTTGG

**Flow Cytometry.** Cells were stained as described.<sup>10,12</sup> Data are representative of >3 experiments.




Figure 2. Gli2 $\Delta$ C2 transgene expression improves TCR-induced activation and proliferation in peripheral T cells. (A) Histograms showing the early activation marker CD69 on CD4 and CD8 SP cells after 24 hours in culture stimulated with 0.01  $\mu$ g/ml of each of anti-CD3 and anti-CD28. Numbers indicate the percentage of cells falling within the marker. Bar chart shows relative increase in activation compared to WT littermate controls. (B) Histograms showing the later activation marker CD25 on CD4 and CD8 SP cells after 24 hours in culture stimulated with 0.01  $\mu$ g/ml of each of anti-CD3 and anti-CD28. Numbers indicate the percentage of cells falling within the marker. Bar chart shows relative increase in activation compared to WT littermate controls. (C) CFSE staining in peripheral T cells cultured for 72 hours with 0.01  $\mu$ g/ml of each of anti-CD3 and anti-CD28. Bar graph to show the number of cell divisions that had occurred. (D) Histograms of CFSE staining to show dose effect of the Gli2 $\Delta$ C2 transgene on proliferation.

**In vitro T cell culture and activation.** Splenic T cells were cultured and activated as described.<sup>12</sup> r-Shh was a gift from Curis.<sup>15</sup>

#### Acknowledgements

This work was supported by the MRC, Leukaemia Research Fund, BBSRC and the Wellcome Trust.

We thank Zoe Webster for microinjection, Hiroshi Sasaki for the Gli $\Delta$ C<sub>2</sub> plasmid, and Curis for the Shh.

#### References

- Crompton T, Outram SV, Hager-Theodorides AL. Sonic hedgehog signalling in T-cell development and activation. *Nature reviews* 2007; 7:726-35.
- Ingham PW, McMahon AP. Hedgehog signaling in animal development: paradigms and principles. *Genes Dev* 2001; 15:3059-87.
- Rowbotham NJ, Hager-Theodorides AL, Furmanski AL, Crompton T. A novel role for Hedgehog in T-cell receptor signaling: implications for development and immunity. *Cell cycle* 2007; 6:2138-42.
- Gill PS, Rosenblum ND. Control of murine kidney development by sonic hedgehog and its GLI effectors. *Cell cycle* 2006; 5:1426-30.
- Park HL, Bai C, Platt KA, Matisse MP, Beeghly A, Hui CC, Nakashima M, Joyner AL. Mouse Gli1 mutants are viable but have defects in SHH signaling in combination with a Gli2 mutation. *Development* 2000; 127:1593-605.
- Sasaki H, Nishizaki Y, Hui C, Nakafuku M, Kondoh H. Regulation of Gli2 and Gli3 activities by an amino-terminal repression domain: implication of Gli2 and Gli3 as primary mediators of Shh signaling. *Development* 1999; 126:3915-24.
- Aza-Blanc P, Lin HY, Ruiz i Altaba A, Kornberg TB. Expression of the vertebrate Gli proteins in *Drosophila* reveals a distribution of activator and repressor activities. *Development* 2000; 127:4293-301.
- Bai CB, Auerbach W, Lee JS, Stephen D, Joyner AL. Gli2, but not Gli1, is required for initial Shh signaling and ectopic activation of the Shh pathway. *Development* 2002; 129:4753-61.
- El Andaloussi A, Graves S, Meng F, Mandal M, Mashayekhi M, Aifantis I. Hedgehog signaling controls thymocyte progenitor homeostasis and differentiation in the thymus. *Nat Immunol* 2006; 7:418-26.
- Hager-Theodorides AL, Dessens JT, Outram SV, Crompton T. The transcription factor Gli3 regulates differentiation of fetal CD4<sup>+</sup> CD8<sup>-</sup> double-negative thymocytes. *Blood* 2005; 106:1296-304.
- Shah DK, Hager-Theodorides AL, Outram SV, Ross SE, Varas A, Crompton T. Reduced thymocyte development in sonic hedgehog knockout embryos. *J Immunol* 2004; 172:2296-306.
- Rowbotham NJ, Hager-Theodorides AL, Cebeacuer M, Shah DK, Drakopoulou E, Dyson J, Outram SV, Crompton T. Activation of the Hedgehog signaling pathway in T-lineage cells inhibits TCR repertoire selection in the thymus and peripheral T-cell activation. *Blood* 2007; 109:3757-66.
- Stewart GA, Lowrey JA, Wakelin SJ, Fitch PM, Lindey S, Dallman MJ, Lamb JR, Howie SE. Sonic hedgehog signaling modulates activation of and cytokine production by human peripheral CD4<sup>+</sup> T cells. *J Immunol* 2002; 169:5451-7.
- Lowrey JA, Stewart GA, Lindey S, Hoyne GF, Dallman MJ, Howie SE, Lamb JR. Sonic hedgehog promotes cell cycle progression in activated peripheral CD4(+) T lymphocytes. *J Immunol* 2002; 169:1869-75.
- Outram SV, Varas A, Pepicelli CV, Crompton T. Hedgehog signaling regulates differentiation from double-negative to double-positive thymocyte. *Immunity* 2000; 13:187-97.
- Sacedon R, Diez B, Nunez V, Hernandez-Lopez C, Gutierrez-Frias C, Cejalvo T, Outram SV, Crompton T, Zapata AG, Vicente A, Varas A. Sonic hedgehog is produced by follicular dendritic cells and protects germinal center B cells from apoptosis. *J Immunol* 2005; 174:1456-61.
- Buckland J, Pennington DJ, Bruno L, Owen MJ. Co-ordination of the expression of the protein tyrosine kinase p56(lck) with the pre-T cell receptor during thymocyte development. *European journal of immunology* 2000; 30:8-18.
- Agathocleous M, Locker M, Harris WA, Perron M. A general role of hedgehog in the regulation of proliferation. *Cell cycle* 2007; 6:156-9.
- Jensen AM, Wallace VA. Expression of Sonic hedgehog and its putative role as a precursor cell mitogen in the developing mouse retina. *Development* 1997; 124:363-71.
- Moshiri A, McGuire CR, Reh TA. Sonic hedgehog regulates proliferation of the retinal ciliary marginal zone in posthatch chicks. *Dev Dyn* 2005; 233:66-75.
- Wang Y, Dakubo GD, Thurig S, Mazerolle CJ, Wallace VA. Retinal ganglion cell-derived sonic hedgehog locally controls proliferation and the timing of RGC development in the embryonic mouse retina. *Development* 2005; 132:5103-13.
- Neumann CJ. Hedgehogs as negative regulators of the cell cycle. *Cell cycle* 2005; 4:1139-40.
- Locker M, Agathocleous M, Amato MA, Parain K, Harris WA, Perron M. Hedgehog signaling and the retina: insights into the mechanisms controlling the proliferative properties of neural precursors. *Genes Dev* 2006; 20:3036-48.
- Chaffin KE, Beals CR, Wilkie TM, Forbush KA, Simon MI, Perlmutter RM. Dissection of thymocyte signaling pathways by in vivo expression of pertussis toxin ADP-ribosyltransferase. *Embo J* 1990; 9:3821-9.
- Kim D, Peng XC, Sun XH. Massive apoptosis of thymocytes in T-cell-deficient Id1 transgenic mice. *Molecular and cellular biology* 1999; 19:8240-53.