

Enhancing cardiometabolic health through physical activity and breakfast manipulations in children and adolescents: good for humans, good for the planet

Julia K Zakrzewski-Fruer¹ & Alice E Thackray^{2,3}

¹Institute for Sport and Physical Activity Research, School of Sport Science and Physical Activity, University of Bedfordshire, MK41 9EA, UK

²National Centre for Sport and Exercise Medicine, School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough LE11 3TU, UK.

³National Institute for Health Research (NIHR) Leicester Biomedical Research Centre, University Hospitals of Leicester NHS Trust and University of Leicester, Leicester LE5 4PW, UK.

Corresponding author: Dr. Julia K Zakrzewski-Fruer, Institute for Sport and Physical Activity Research, School of Sport Science and Physical Activity, University of Bedfordshire, Bedford, MK41 9EA, UK. Telephone: +44 (0)1234 793410. E-mail: Julia.Fruer@beds.ac.uk

Short title: Enhancing metabolic health in young people

Key words: Climate change, diabetes, disease prevention, exercise, youth

Acknowledgements

AET acknowledges support from the National Institute for Health Research (NIHR) Leicester Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health.

Financial Support

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest

None.

Authorship

The authors had sole responsibility for all aspects of preparation of this paper.

Abstract

The human health benefits of cardiometabolic disease prevention can be accompanied by planetary co-benefits. Focusing efforts towards young people, including children and adolescents, is conducive to disease prevention. In the context of cardiometabolic disease prevention, this review paper critically summarises the available literature on the acute cardiometabolic responses to physical activity and breakfast manipulations among young people. Given the seriousness of global climate change, which will disproportionately affect our younger generations, the paper offers new insights into the inherent interactions between child-adolescent behaviour and cardiometabolic health from an environmental sustainability perspective to aid climate change mitigation efforts, including exploring future research avenues. A growing evidence base suggests acute moderate- to high-intensity exercise bouts can attenuate postprandial plasma glucose, insulin and triacylglycerol concentrations for up to 24–48 h in young people. Whether accumulating physical activity throughout the day with short, frequent bouts promotes cardiometabolic risk marker attenuations is unclear. Breakfast consumption may enhance free-living physical activity and reduce glycaemic responses to subsequent meals for a possible additive impact. If repeated habitually, attenuations in these cardiometabolic risk factors would be conducive to disease prevention, reducing the greenhouse gas emissions associated with disease diagnosis and treatment. To progress current understanding with high public health and planetary relevance, research among samples of ‘at risk’ young people that span cellular-level responses to ecologically valid settings and address human and planetary health co-benefits is needed. Indeed, certain physical activity opportunities, such as active travel to school, offer important direct co-benefits to human and planetary health.

Introduction: childhood and adolescence as ‘critical periods’

The prevalence of cardiometabolic disease risk markers, including insulin resistance and dyslipidaemia, and even the manifestations of certain cardiometabolic diseases are increasing globally among young people⁽¹⁻³⁾. Given that cardiometabolic disease develops over time, an understanding of disease precursors rather than manifestation is more relevant to young people. Even so, in the absence of global data, recent data of 3.47 million 10–19-year-olds in the U.S reported a 95.3% relative increase in estimated type 2 diabetes prevalence over 16 years between 2001 and 2017⁽³⁾. Furthermore, U.S National Health and Nutrition Examination Survey data has shown that 18.0% of 12–18-year-olds have prediabetes⁽¹⁾ and 20.2% of 8–17-year-olds have an adverse lipid concentration⁽⁴⁾. With rises in obesity among young people being a key driver, this data suggests cardiometabolic disease is set to manifest at an earlier age in future generations, prolonging the health and economic burden^(1,5). Importantly, targeting young people to aid disease prevention rather than focusing efforts toward treatment in later life may also benefit the planet through reduced greenhouse gas emissions associated with disease diagnosis and treatment within healthcare systems; for example due to reduced manufacturing of blood analysis consumables and drugs, motorised transportation of goods and services, and clinical waste disposal (see Fig. 1)⁽⁶⁾.

Moderating postprandial glycaemia, insulinemia, and lipaemia is at the cornerstone of cardiometabolic disease prevention^(7,8). Such risk markers share common physiological pathways, leading to their ‘clustering’ and placing young people at further risk⁽⁵⁾. As such, public health recommendations that address multiple cardiometabolic disease risk markers stand to create the largest impact. Most waking hours are spent inducing marked metabolic fluctuations, emphasising the importance of postprandial rather than fasting measures. Postprandial glycaemia can increase cardiometabolic disease risk through oxidative stress, inflammation, and atherosclerosis^(7,8) with repeated rises and falls indicative of high glycaemic variability appearing particularly harmful^(9,10). Even in ‘healthy’ populations, higher postprandial glucose concentrations increase type 2 diabetes, cardiovascular disease and all-cause mortality risk⁽¹¹⁻¹⁴⁾. Likewise, regular exposure to elevated postprandial triacylglycerol concentrations promotes a pro-atherogenic lipid phenotype, including low HDL concentrations and the accumulation of small, dense LDL⁽¹⁵⁾ that increases susceptibility to future clinical cardiovascular events. Taken together, these findings may explain the lack of sensitivity of fasting measures alone for the diagnosis of cardiometabolic disease⁽¹⁶⁾.

Childhood and adolescence are vulnerable times of life and have been associated with entering the initial stages of cardiometabolic disease development. Among girls and boys, pubertal insulin resistance reduces the ability of insulin to increase glucose uptake and utilisation, resulting in a

compensatory increase in insulin secretion and plasma insulin concentrations to maintain normal blood glucose concentrations^(17,18). Pubertal insulin resistance is transient, typically reported to begin during early puberty, peak during mid-puberty and recover by the end of puberty⁽¹⁷⁾. Yet, more recent longitudinal data has demonstrated that the first stages are detectable from ~7 years of age⁽¹⁹⁾. Unfortunately, the diets of many young people are not conducive to the puberty-related reduction in insulin sensitivity; for example, free sugar intakes exceeded the recommended level in 87% of 4–10-year-olds and 92% of 11–18-year-olds in the UK in 2020⁽²⁰⁾. The factors contributing to pubertal insulin resistance remain elusive, but appear to include rises in insulin-like growth factor-1 and adiposity^(19,21,22), and affect girls more than boys^(18,19,23). Due to the additive effect of adiposity, unfortunately pubertal insulin resistance in young people with obesity does not fully resolve, resulting in an increased risk of β -cell exhaustion and cardiometabolic disease at a younger age⁽²¹⁾. Likewise, the antecedents of atherosclerotic disease originate in the formative years⁽²⁴⁾, where aortic and coronary atherosclerotic lesions can develop in the first two decades of life and progress in prevalence and severity over time^(25,26). Further, cardiovascular disease risk markers that manifest in youth are predictive of subclinical atherosclerosis in adulthood^(27,28) and may increase the risk of premature death in adults⁽²⁹⁾. This data highlights the importance of protecting cardiometabolic health from an early age.

For effective cardiometabolic disease prevention, childhood and adolescence have been identified as a critical time to establish healthy, sustainable physical activity and dietary behaviours for life-long impact^(30,31). Yet, young people remain under-represented in the field of exercise and nutrient metabolism. Drawing on data from adult samples, acute exercise is a potent stimulus for moderating postprandial glycaemia, insulinaemia and lipaemia for up to two to three days^(32,33,34). Critically, such findings may not apply to the distinct hormonal, metabolic and behavioural profiles of children and adolescents, including pubertal insulin resistance^(17,18), fuel utilisation^(35,36) and physical activity patterns⁽³⁷⁾. The lack of paediatric-specific data is concerning given the low physical activity levels of many young people; for example, ~71% of European 2–18-year-olds do not meet the physical activity guidelines when measured objectively⁽³⁸⁾, and ~81% of 11–17-year-olds were classified as insufficiently physically active globally based on population survey data of 1.6 million participants⁽³⁹⁾. Attention to adolescents is of particular relevance due the adolescent decline in physical activity, which is more pronounced among girls than boys and supported by a large body of longitudinal research employing objective physical activity measures⁽⁴⁰⁾. During this time, adolescents often begin to skip breakfast⁽⁴¹⁾, which may interact with physical activity and glycaemic control^(42–48). However, a comprehensive summary of the available paediatric evidence

on the acute glycaemic, insulinaemic and lipaemic responses to physical activity and the role of breakfast-physical activity interactions has not yet been undertaken.

This review paper will critically summarise the extant literature on the cardiometabolic responses up to 48 h after completing physical activity bouts among children and adolescents. The possible interactions of breakfast with physical activity on cardiometabolic disease risk are then discussed. Reflecting traditional cardiometabolic disease risk markers and the available paediatric literature, this review paper will pay specific attention to glycaemia, insulinaemia and lipaemia, notwithstanding that an array of other markers exists. Given the seriousness and timeliness of global climate change with responsibility to act across disciplines, this review paper offers new insights into the inherent interactions between child-adolescent behaviour and cardiometabolic health in the context of environmental sustainability to aid climate change mitigation efforts, including exploring future research avenues. Based on these themes, Fig. 1 provides a platform for the concepts integral to this review paper, offering potential avenues that support a focus on young people for cardiometabolic disease prevention with co-benefits to mitigating climate change through direct and indirect pathways. From the outset, it should be highlighted that the term ‘physical activity’ is used to describe bodily movements with an energy expenditure >1.5 METs, whereas the term ‘exercise’ is used to describe the subset of physical activity that has pre-planned, structured characteristics. The collective term ‘young people’ is used where findings apply to mixed samples of children and adolescents and where it is not essential to distinguish between chronological age (referred to as ‘age’ for simplicity) and pubertal stage.

<Please insert Fig. 1 near here>

Cardiometabolic responses to acute physical activity

As illustrated in Fig. 2A, experimental investigations on acute physical activity have typically employed crossover designs using a one- or two-day study paradigm, whereby the physical activity stimulus is either applied before measuring fasting and/or postprandial cardiometabolic responses or is accumulated throughout the measurement period. These experimental designs over multiple days reflect the expected lag between exposure to the physical activity stimulus and cardiometabolic benefit. The test day protocols for the measurement of outcome variables are designed in line with the differing pattern, duration and magnitude of postprandial glycaemic, insulinaemic and lipaemic responses to a single or multiple meals consumed across the day, as shown in Fig. 2B.

<Please insert Fig. 2 near here>

Single exercise bouts

Glycaemia and insulinaemia

With children and adolescents being under-represented in the field of exercise metabolism, research on the glycaemic and insulinaemic responses to acute exercise in paediatric populations began to emerge around 10 to 15 years after^(49–52) the adult-based literature was reviewed⁽³²⁾. To date, studies in samples of healthy adolescent boys have shown that acute moderate-intensity exercise and high-intensity interval exercise (HIIE) can improve indices of glucose control and insulin sensitivity assessed during an oral glucose tolerance test (OGTT) immediately⁽⁵⁰⁾ and for up to 24 h⁽⁵¹⁾ compared with no exercise. In contrast, 7–10-year-old boys had limited scope for improvement in OGTT-derived indices of glucose control and insulin sensitivity after single bouts of HIIE and moderate-intensity exercise, perhaps due to their younger age⁽⁵²⁾. The OGTT is primarily intended for diabetes diagnosis and does not necessarily reflect the glycaemic challenges of habitual dietary intakes. Using more ‘ecologically valid’ test meals among female and male healthy adolescents, 60-min of games-based activity improved insulin sensitivity in response to a standardised lunch consumed 60-min post-exercise⁽⁵³⁾. However, a 60-min school-based football session was only sufficient to reduce glucose concentrations 60-min post-exercise and did not affect the total glucose response (i.e., the area under the curve) or the insulin response⁽⁵⁴⁾. Thus, the glycaemic and insulinaemic benefits of acute exercise among young people reported in controlled laboratory environments may not necessarily translate to more ecologically valid settings.

The persistence of improvements in glycaemia and insulinaemia in the days after exercise versus rest has been examined in adolescents described as ‘healthy’ and ‘at risk’ of cardiometabolic disease. Beneficial effects in indices of insulin sensitivity and glucose control were reported 17-h after a 45-min aerobic exercise bout at 75% peak heart rate in response to a standardised mixed meal among female and male adolescents with habitually low physical activity levels⁽⁵⁵⁾ and after moderate-intensity exercise and HIIE in response to an OGTT 24-h post-exercise among healthy adolescent boys⁽⁵¹⁾. Such effects may interact with participant characteristics, including weight status; indeed, a bout of exercise at Fatmax (i.e., the intensity that elicits individual peak fat oxidation) with an estimated energy expenditure of 2092 kJ performed 16-h before a high-glycaemic-index breakfast reduced fasting and postprandial insulin concentrations in girls classified as non-overweight (i.e., ‘healthy’ weight), but not in girls classified as overweight/obese⁽⁴⁹⁾. The lack of change in the girls with overweight/obesity may reflect a degree of metabolic inflexibility or a higher energy and carbohydrate intake to compensate for the exercise-induced energy deficit in this group⁽⁴⁹⁾. In healthy mixed-sex or male-only samples of adolescents, no effects of 30- or 60-

min of high-intensity intermittent running were seen 24-h post-exercise in response to an ecologically valid breakfast meal⁽⁵⁶⁾ or at 24- or 48-h after moderate-intensity, HIIE or games-based activity using fasted measures^(51,53). Thus, the time-course of potential post-exercise improvements in glycaemia and insulinaemia among young people requires further investigation, taking into account participant characteristics.

A higher exercise intensity and/or longer duration may enhance the potential for exercise-induced improvements in glucose control and insulin sensitivity due to increased glycogen depletion providing a stimulus for muscle glycogen repletion via GLUT-4, as suggested by adult data⁽³²⁾. In line with this, more pronounced glycaemic and insulinaemic benefits among adolescents were seen in response to a 60- versus 30-min bout of high-intensity intermittent running in girls and boys⁽⁵⁶⁾ and in response to work-matched high-intensity interval cycling versus moderate-intensity exercise in boys⁽⁵¹⁾. Given the importance of muscle glycogen depletion, it should be highlighted that the nutritional status of participants varies between studies, with exercise performed in the fed^(49,54,56) or fasted^(50,51) state. Although dietary intakes on the day of the exercise intervention can often be controlled, this is challenging when participants are unsupervised by the researchers. Thus, control of dietary intake between experimental trials has often been implemented within, but not between participants, using replication of habitual intakes via food diaries^(49,51). Such issues may contribute to conflicting findings across studies.

Related to the exercise-induced energy deficit and muscle glycogen depletion, enhanced fat oxidation may partly underpin improvements in insulin sensitivity after acute exercise through a reduced accumulation of fatty acid metabolites (e.g., diacylglycerol, ceramides) within the muscle that can interfere with insulin signalling^(57,58). Such mechanisms could have high relevance to young people with obesity who can already exhibit markers of fatty acid-induced insulin resistance⁽⁵⁹⁾. Although studies that have measured fatty acid metabolite accumulation in response to exercise in young people are lacking, increased fat oxidation rates have been reported immediately after moderate-intensity exercise and HIIE in adolescent boys⁽⁵⁰⁾ and ~16-h after exercise among adolescent girls classified as overweight/obese or non-overweight⁽⁴⁹⁾, whereas 7–10-year-old boys exhibited an increase in fat oxidation immediately after HIIE but not moderate-intensity exercise⁽⁵²⁾. It possible that such effects may not apply equally across different age groups and pubertal stages because the child-adolescent-adult transition is accompanied by a reduced reliance on fat and exogenous carbohydrate with a concomitant increased reliance on total and endogenous carbohydrate as fuels during exercise, which is governed by advances in puberty rather than age^(35,36). Yet, studies have not typically controlled for pubertal stage, often due to ethical issues

associated with the recruitment of participants based on this sensitive marker. Although girls may have higher glucose and insulin responses than boys during puberty, they appear to respond similarly to acute exercise^(53,56); thus, the influence of sex on between-study comparisons may be minimal.

Although important paediatric literature has accumulated over recent years, an understanding of exercise-induced improvements in glycaemia and insulinaemia remains in its infancy. This data suggests that improvements in glucose control and insulin sensitivity can be achieved through acute exercise bouts. Yet, the extent of such improvements appears to depend on the exercise characteristics (e.g., intensity, duration and mode), sample characteristics (e.g., pubertal stage, sex, baseline insulin resistance, habitual physical activity levels, adiposity), and possible compensatory physical activity and dietary intake responses during the often uncontrolled period between the intervention and outcome measures. Such factors require further investigation to inform physical activity recommendations. With much of the existing paediatric literature based on healthy young people, further research in samples of girls and boys who are at risk of cardiometabolic disease would have valuable public health impact. An insight into the underpinning physiological mechanisms is also required. Indeed, different mechanisms may be at play when compared with adults due to the limited glycogen storage capacity and lower reliance on endogenous glycogen as a fuel in younger, less mature children and adolescents^(35,36,60). That said, it may be possible to apply findings based on adults more readily to adolescents in the later stages of puberty once an ‘adult-like’ metabolic profile has been developed⁽³⁵⁾ and after exercise training in boys, where an increased capacity to store and use muscle glycogen has been shown⁽⁶⁰⁾. Yet, since the early muscle biopsy studies in the 1970s and 1980s⁽⁶¹⁾, the invasive measures often required to provide insight at the cellular level have been bound to more stringent ethical and technical constraints than the adult literature. More promisingly, advances in technologies have enabled minimally invasive techniques to be employed with young people, including stable isotope tracer techniques and magnetic resonance spectroscopy^(35,36,59).

Lipaemia

Scientific interest exploring postprandial lipaemic responses to exercise in young people has grown in the last two decades. The maximal triacylglycerol-lowering effect of acute exercise is typically delayed until 12–20-h after exercise, coinciding with the exercise-mediated increase in lipoprotein lipase activity that facilitates circulating triacylglycerol clearance⁽³⁴⁾. Consequently, most paediatric studies have adopted a two-day model whereby the exercise stimulus is applied the day before measuring metabolic responses to high-fat meals (Fig. 2A). This body of work has shown

consistently that moderate- to high-intensity exercise bouts performed 12–18-h before a high-fat meal reduces postprandial triacylglycerol concentrations. Although the magnitude of reduction varies highly across studies, the TAG-lowering effect of acute exercise has been demonstrated in both boys and girls classified as lean, overweight and moderately obese. In contrast, using a one-day protocol, moderate- and high-intensity exercise bouts have yielded negligible effects^(62,63). Along with increased lipoprotein lipase activity, the secretion of fewer, triacylglycerol-rich, very low-density lipoprotein particles from the liver that have a higher affinity for lipoprotein lipase have been proposed in adults as a central mechanistic pathway responsible for the exercise-induced reduction in postprandial lipaemia⁽⁶⁴⁾.

A key determinant of the reduction in postprandial lipaemia the day after exercise in adults is the exercise-induced energy expenditure^(33,34). In contrast, there is little evidence of clear dose-dependent relationships between exercise energy expenditure and reduced postprandial triacylglycerol concentrations in young people^(65–67). Specifically, similar exercise-induced reductions in postprandial lipaemia were observed in 12–14-year-old boys when comparing two exercise bouts varying in intensity (60-min at 53 versus 75% of peak oxygen uptake)⁽⁶⁵⁾ and duration (30- versus 60-min at 55% of peak oxygen uptake)⁽⁶⁶⁾ despite a two-fold difference in gross energy expenditure. Furthermore, a 60-, but not 30-min, exercise bout (1536 versus 777 kJ, respectively) reduced postprandial triacylglycerol concentrations in 10–14-year-old girls, and does not support a dose-dependent relationship given the 30-min bout did not induce an ‘intermediate’ effect⁽⁶⁷⁾. Nevertheless, ~56% of the female participants experienced a reduction in postprandial triacylglycerol concentrations after 30-min of exercise, suggesting some girls may accrue metabolic health benefits from lower volumes of exercise⁽⁶⁷⁾.

The triacylglycerol-lowering effects of single exercise bouts have been demonstrated consistently in response to laboratory-based exercise protocols involving moderate- to vigorous-intensity treadmill walking or running and cycling that meet the current physical activity guidelines for young people^(65–74). Alternatively, other studies have also highlighted the efficacy of acute intermittent^(68,75) or high-intensity^(76–79) exercise regimens for reducing postprandial lipaemia early in life. Such protocols may better match the sporadic stop-start activity habits of young people⁽³⁷⁾ and offer an alternative form of activity that can evoke important metabolic health benefits. Desirable reductions in postprandial lipaemia have been reported in response to repeated 1-min running intervals performed at 100% of peak oxygen uptake^(76,77) in addition to repeated short (~6-s) maximal running⁽⁷⁸⁾ and cycling⁽⁷⁹⁾ sprints. In the earliest exercise postprandial study in young people, simulated games activity comprising repeated blocks of walking, sprinting, cruising,

jogging and rest reduced postprandial lipaemia by 26% in adolescent boys (mean age 15.4 years)⁽⁶⁸⁾. A similar reduction in postprandial triacylglycerol concentrations was reported more recently the day after 11–13-year-old boys completed a 48-min session of school-based free-living soccer⁽⁷⁵⁾. The magnitude of reduction after the soccer activity was comparable to that observed after time-matched moderate-intensity treadmill exercise (25% versus 18%, respectively), but importantly provides a more representative form of activity for young people and advances insights gained from laboratory-based studies that dominate the paediatric literature⁽⁷⁵⁾.

Beneficial effects of single exercise bouts on postprandial lipaemia have been observed in boys and girls classified as overweight or moderately obese^(69,70). A single bout of 60-min treadmill exercise at 65% of peak oxygen uptake evoked an equivalent reduction (~20%) in postprandial lipaemia in late-adolescent boys classified as lean or overweight⁽⁶⁹⁾. More recently, 60-min of moderate-intensity cycling reduced postprandial triacylglycerol concentrations in black and white adolescents (12–18-years-old) classified as overweight⁽⁷⁰⁾. The triacylglycerol-lowering effects of exercise were more pronounced in the white than black adolescents (19% versus 8%, respectively), with increased visceral adipose tissue identified as a strong determinant of the exercise-induced reduction in postprandial lipaemia in the white adolescents⁽⁷⁰⁾. Despite the elevated adiposity status, the participants in these studies were otherwise healthy. Nevertheless, these findings provide a foundation for future work in young populations at elevated cardiometabolic disease risk (e.g., hypertriglyceridaemia, insulin resistant) where augmenting postprandial metabolic health may have most clinical relevance.

Another avenue of inquiry has investigated whether the acute benefit of exercise on postprandial lipaemia is due to the associated energy deficit or driven by skeletal muscle contraction itself. Immediate dietary replacement of the exercise energy expenditure counteracted the reduction in postprandial triacylglycerol concentrations observed the day after a bout of moderate-intensity exercise in 11–13-year-old boys⁽⁷⁴⁾. Thus, maintenance of the energy deficit in the immediate post-exercise period may be required to maximise the metabolic health benefits. Other studies have explored the potential differential effects of imposing energy deficits from acute exercise and/or energy intake restriction. This work has demonstrated that a moderate-intensity exercise (~1.5 MJ) bout was a more potent stimulus for reducing postprandial lipaemia than an equivalent diet-induced energy deficit in 11–13-year-old girls⁽⁷³⁾. Subsequently, Thackray et al.⁽⁷⁷⁾ compared the effect of 10 × 1-min high-intensity treadmill runs and 5 × 1-min high-intensity treadmill runs combined with a controlled restriction in habitual energy intake at the evening meal (~0.82 MJ) in girls (mean age 12.1 years). The reduction in postprandial triacylglycerol concentrations reported after high-

intensity running alone was similar in magnitude when the smaller exercise dose was superimposed with energy intake restriction (10% versus 9% respectively)⁽⁷⁷⁾. Therefore, combinations of exercise and mild dietary energy intake restriction may offer a practical alternative for young people to acquire metabolic health benefits, particularly those who struggle to accumulate a sufficient level of physical activity.

Short, frequent physical activity bouts

Short, frequent physical activity bouts that break up sedentary behaviour have received growing interest as evidence of the high prevalence of sedentary behaviour across the lifespan continues to accumulate⁽⁸⁰⁾. Sedentary behaviours are any waking behaviours characterised by an energy expenditure ≤ 1.5 METs, while in a sitting, reclining, or lying posture⁽⁸¹⁾, which can be replaced by physical activity of a light (1.6-2.9 METs), moderate (3.0-5.9 METs) or vigorous (≥ 6.0 METs) intensity during waking hours. This avenue of research is timely as sedentary behaviour has emerged as a strong determinant of mortality and cardiometabolic disease risk in adults^(82,83) and is associated with adverse cardiometabolic health outcomes in young people⁽⁸⁴⁾. Efforts to mitigate the deleterious effects of prolonged sitting (a common proxy for sedentary behaviour) through short, frequent physical activity bouts are likely to have greater appeal for those who are not motivated to perform continuous bouts of structured exercise, where paediatric populations typically perceive prolonged exercise as more demanding than adults⁽⁸⁵⁾. This pattern of activity may also more closely reflect the sporadic physical activity patterns of children and adolescents⁽³⁷⁾ and be more conducive to integrate into the school day.

A meta-analysis of studies across the lifespan showed that interrupting sitting with short, frequent light- to moderate-intensity physical activity bouts reduces postprandial glucose and insulin, but not triacylglycerol, concentrations⁽⁸⁶⁾. Although the impact of disrupting sitting with regular activity bouts on postprandial cardiometabolic risk markers was not modified by age, a closer examination of the paediatric-specific studies presented here suggests relatively modest and inconsistent effects. In this regard, breaking up prolonged sitting with 3-min moderate-intensity walking bouts performed every 30-min during a 3-h OGTT reduced glucose and insulin concentrations but did not alter postprandial lipaemia in 7–11-year-old children with healthy weight⁽⁸⁷⁾ and reduced insulin concentrations in 7–11-year-old children who were overweight or obese⁽⁸⁸⁾. Using less frequent physical activity bouts in 13–14-year-old boys, accumulating four bouts of work-matched high-intensity physical activity (2×1 -min intervals at 90% peak power) reduced the postprandial glucose response on the intervention day when compared with continuous moderate-intensity (~ 6 -min at 90% gas exchange threshold) exercise and a rest control trial, although no effect was

observed for postprandial lipaemia⁽⁸⁹⁾. In contrast, postprandial glucose, insulin and triacylglycerol concentrations were all unaffected in response to interrupting sitting with 2-min light-intensity walking bouts every 20-min for 8-h compared to 8-h of uninterrupted sitting in 10–14-year-olds; this lack of effect was maintained even when the 2-min light walking breaks were supplemented with two, 20-min moderate-intensity treadmill bouts⁽⁹⁰⁾. Similarly, breaking up sitting time with bouts of self-paced walking and standing accumulated during two consecutive simulated school days did not alter postprandial glucose, insulin or triacylglycerol concentrations assessed on the second simulated school day or the next day in 11–13-year-old physically inactive girls⁽⁹¹⁾. Thus, the immediate and delayed effect of an ecologically valid standing and walking intervention appears to be negligible.

Discrepancies between studies are most likely explained by differences in participant characteristics and study design features, including variations in the physical activity or exercise stimulus, meal content or glycaemic load, timing of activity bouts and frequency of outcome assessments. Specifically, the relatively low frequency of blood sampling of ≥ 90 -min intervals in some studies may have missed important glycaemic fluctuations in close proximity to the meals and activity/standing bouts^(90,91). Using continuous glucose monitoring devices that measure interstitial glucose concentrations at ~ 5 -min intervals, interrupting prolonged sitting with 2-min resistance activity bouts every 18-min reduced the incremental, but not total, glucose area under the curve for 3-h after both breakfast and lunch meals in adolescents aged 14–17-years⁽⁹²⁾. Yet, these changes were relatively subtle in magnitude with detection likely facilitated by the improved sensitivity of continuous glucose monitors to capture glucose dynamics.

The lack of consistent reductions in postprandial lipaemic responses may be a consequence of the 1-day study protocol often employed in these studies^(87–90). In this regard, accumulating 6×10 -min bouts of running at 70% of peak oxygen uptake did not alter next day postprandial glucose and insulin concentrations, but did reduce total and incremental area under the triacylglycerol curve by 11% and 16%, respectively, in adolescent boys aged 12–14 years⁽⁷²⁾. These differences did not reach statistical significance, but are similar in magnitude to other exercise postprandial studies in children and adolescents. In support of the delayed effect of exercise on postprandial lipaemia, an in-school protocol in 11–13-year-old boys comparing the effect of 40×30 m sprints in one uninterrupted after school bout and accumulated in four bouts of 10×30 m sprints strategically placed during natural breaks in the school day reported reductions in postprandial triacylglycerol concentrations the next day, despite no effect on the day of exercise⁽⁷⁸⁾. This may provide a viable model for translating the metabolic health benefits reported with laboratory-based exercise

protocols into free-living settings providing the exercise stimulus is of sufficient duration and/or intensity. Importantly, since favourable effects among adults are most consistently seen in clinical, overweight/obese, and physically inactive populations⁽⁸⁶⁾, further research in such paediatric populations is warranted and has high public health relevance.

Acute breakfast consumption-physical activity interactions for cardiometabolic health

Cross-sectional and prospective observational studies show that frequent breakfast consumption is associated with reduced cardiometabolic disease risk in young people^(41,93) and manifestation in adults^(94,95). Such findings may in part be explained by higher free-living physical activity and enhanced glucose control in response to breakfast consumption versus breakfast omission, as reported in randomised controlled trials and crossover trials in adults^(42–47). An eating pattern that includes breakfast to aid the provision of a continuous carbohydrate supply may be of greater importance to young people versus adults due to their elevated resting and physical activity energy expenditures^(96,97) and higher reliance on exogenous carbohydrate as a fuel⁽³⁶⁾. Indeed, increased exogenous glucose availability (i.e., a physiological response) as well as perceived energy levels (i.e., a psychological response) are potential mediators of the breakfast-physical activity relationship. An understanding of such responses is particularly relevant to adolescent girls due to the adolescent decline in breakfast consumption frequency and physical activity levels as well as pubertal insulin resistance disproportionately affecting girls^(18,19,23,40,41). Depending on the country and definition of breakfast employed, the prevalence of skipping breakfast can range from 10–30% among young people, and is higher among adolescents (versus children) and girls (versus boys)⁽⁴¹⁾.

Inconsistent cross-sectional findings on the association between breakfast consumption and objectively-measured physical activity among young people^(98–100) highlight the need for experimental trials to provide more definitive causative interpretations of the findings with reduced interference from confounders. Using a 3-day crossover trial to compare standardised breakfast consumption versus breakfast omission in adolescent girls, physical activity was unaffected despite incomplete energy intake compensation of the breakfast meal⁽¹⁰¹⁾. If repeated habitually, this lack of energy compensation would be expected to result in weight gain in response to breakfast consumption versus omission, contradicting the higher adiposity reported among habitual breakfast skippers^(41,93). Yet, it is possible that wrist-worn accelerometry used in this study was not sensitive enough to detect differences in physical activity in response to breakfast manipulation. Using combined heart rate accelerometry, a 7-day crossover trial reported that adolescent girls spent more time in light-intensity physical activity before 10:30 and after school, and less time sedentary after school during daily versus intermittent (i.e., alternate breakfast omission and consumption)

standardised breakfast consumption⁽¹⁰²⁾. The 19.8 min/day higher light-intensity physical activity during daily versus intermittent breakfast consumption complemented previous findings among adults that breakfast-induced changes in physical activity may be characterised by the replacement of sedentary behaviours with spontaneous light-intensity physical activity, whereas moderate- to vigorous-intensity physical activity is typically planned and structured⁽⁴²⁾. Interestingly, no differences were reported during the 10:30–15:00 time segment, possibly because the school timetable restricts physical activity opportunities, limiting the ‘window of opportunity’ among school-aged young people. Thus, previous reports that morning physical activity typically before around 12:00 has been shown to be particularly sensitive to breakfast manipulation in adults^(42–44) may not apply to adolescents during school days. Although it is plausible that contrasting the extremes of daily breakfast omission versus consumption across seven days may confer more pronounced effects, no difference in physical activity quantified via combined heart-rate accelerometry was found when comparing these extremes in adolescent girls who habitually skipped breakfast⁽¹⁰³⁾. Nevertheless, the girls were able to accurately compensate after 10:30 for the missed energy at breakfast through increased energy intake, primarily in the form of carbohydrate, during breakfast omission versus breakfast consumption. The lack of physical activity response may have been because the girls were already accustomed metabolically and behaviourally to skipping breakfast, suggesting that habituation to breakfast omission could alter responses to breakfast manipulation. However, ethical restrictions pose challenges in asking girls who habitually consume breakfast to engage in extended periods of breakfast omission.

Inconsistent extant findings may be due to environmental constraints inherent to the lives of adolescent girls, including the school timetable, extra-curricular activities, family routines, and commitments. As such, further research that includes opportunities to be physically active would be useful in extending current knowledge. Adding a further degree of complexity, the nature of the causal relationship between breakfast and physical activity may also depend on physical activity intentions. Indeed, breakfast may promote physical activity or, alternatively, intentions to engage in physical activity may prompt breakfast consumption in preparation, and early morning exercise may result in ‘breakfast consumption’ (i.e., within 2–3-h of waking)⁽¹⁰⁴⁾. Studies in this area have used carbohydrate-based breakfasts to examine the potential impact on free-living physical activity as this is the primary fuel for physical activity^(101–103); nevertheless, the effect of breakfast quantity and quality requires investigation. Additionally, data among boys, younger children and ‘at risk’ young people are required to inform population-specific recommendations.

Due to the lack of clarity on the potential association and/or causal link between breakfast consumption and physical activity among adolescents, alternative mediating factors may play a more prominent role in explaining the favourable cardiometabolic health profiles of habitual breakfast consumers. Among adults, breakfast consumption attenuates glycaemic and insulinaemic responses to subsequent meals when compared to breakfast omission, as termed the ‘second-meal effect’^(45–47). If repeated habitually, a series of moderated glycaemic responses induced by breakfast consumption could reduce cardiometabolic disease risk through a reduced magnitude of glycaemic excursions and glycaemic variability^(7,8). A key mechanism for the ‘second-meal effect’ in adults is increased glucose conversion into muscle glycogen⁽⁴⁷⁾. Such responses may differ in adolescents due to their lower reliance on endogenous glucose and reduced capacity for muscle glycogen storage, particularly in younger, less mature individuals^(35,36,105), and with an estimated 32% reduction in insulin sensitivity occurring from pre- to mid-puberty⁽¹⁷⁾. The slower rate of gastric emptying that has also been proposed to contribute to attenuations in glycaemia in response to breakfast consumption versus omission in adults⁽⁴⁸⁾ may, again, not apply equally to young people. Indeed, puberty, age and sex may affect gastric emptying in healthy children and adolescents, with slower rates reported in girls versus boys and during mid-late puberty versus pre-puberty⁽¹⁰⁶⁾. Unfortunately, current knowledge on breakfast consumption and glycaemia among adolescents relies on a few novel studies that have pooled data from adolescents and adults aged 13–20-year-olds with little consideration of pubertal status, age and sex^(107,108); hence, the responses of children and adolescents specifically remain unknown.

The available research based on pooled adolescent-adult data suggests that the ‘second-meal’ effect depends on factors such as habitual breakfast patterns and breakfast meal composition. In a novel study of girls classified as overweight/obese (mean age 19 years), three days of skipping breakfast versus consuming a normal-protein (12 g protein) breakfast versus consuming a high-protein (32 g protein) breakfast did not affect post-lunch glycaemic and insulinaemic responses when assessed on day 4 among habitual breakfast skippers⁽¹⁰⁷⁾. However, the habitual breakfast consumers, who did not complete the breakfast skipping trial, had lower afternoon and daily glucose responses following the higher-protein breakfast versus the normal-protein breakfast⁽¹⁰⁷⁾. Extending a similar protocol over 12 weeks with a similar yet mixed-sex sample, high protein (35 g) breakfast consumption appeared to be more beneficial for free-living glycaemia than normal protein (13 g) breakfast consumption⁽¹⁰⁸⁾; yet, this was not an acute, controlled study and reductions in energy intake, unhealthy evening snacking of high fat/high sugar foods may have contributed to the findings⁽¹⁰⁹⁾. Other than age and/or puberty, a potential reason for the unclear findings to date may be that the repeated ingestion of glucose appears to be key to inducing the ‘second meal effect’

among adults, with low glycaemic index breakfasts being particularly beneficial^(110,111). Further, the energy content of the breakfast meals^(107,108) was modest when considering data in adults⁽⁴⁵⁻⁴⁷⁾ and recommendations that breakfast should provide at least 20% of daily energy needs^(112,113). Considering daily glucose profiles, low glycaemic index breakfasts also induce a lower immediate response compared with high glycaemic index breakfast among adolescents^(23,114). Using a standardised carbohydrate-rich low glycaemic index breakfast, our recent work with adolescent girls who consumed breakfast habitually demonstrated significant and large reductions in peak, post-lunch (2-h) and total trial (5-h) glycaemia and insulinaemia in response to breakfast consumption versus omission followed by a standardised high glycaemic lunch provided at 3-h, with minimal inter-individual variability⁽¹¹⁵⁾. Further, enjoyment during exercise performed in the afternoon was enhanced with breakfast consumption versus omission⁽¹¹⁵⁾. Although this work did not target adolescents ‘at risk’ of poor glycaemic control, it suggests that moderations in glycaemia with breakfast consumption may be important in driving a favourable cardiometabolic health profile. Further research on the possible interactive effect of breakfast consumption and physical activity requires investigation due to the potential link between these two variables^(98-100,102). Such experimental research would help move closer to developing an evidence-based standardised definition of breakfast consumption for health that can be employed by researchers and practitioners.

Future directions: opportunities to impact human and the planetary health using active travel to school as an example

With preventable cardiometabolic diseases and climate change progressively escalating over recent decades due to a complex interplay of modern human practices, multi-disciplinary approaches that target certain human behaviours may contribute to addressing these global challenges simultaneously. Results on the co-benefits of physical activity and nutrition in young populations for human and planetary health is of high interest, particularly due to cardiometabolic disease risk reaching concerning levels among young people and global climate change placing uncertainty on their futures (see Fig. 1)⁽¹¹⁶⁾. Active travel to school offers a physical activity opportunity for school-aged young people that can benefit both (their) human and planetary health. Further, morning active travel is in close proximity to the breakfast meal, providing a pathway for potential breakfast-physical activity interactions, as discussed previously^(98-100,102). Concurrently, active travel reduces the greenhouse gas emissions associated with motorised vehicle transport, thus assisting climate change mitigation. In contrast to many physical activity opportunities, transport is most often a stipulation of everyday life with almost all school-aged young people travelling to school across the school year. Further, active travel may offer a convenient route to build physical

activity into the day with a minimal time burden and interference with daily tasks; this is important given that lack of time or competing activities for time are common barriers to physical activity⁽¹¹⁷⁾. Thus, encouraging school-based active travel may be a feasible approach to increase population-level physical activity levels.

Accumulating evidence among children and adolescents outlined in this review paper suggests that the likelihood of achieving cardiometabolic health benefits through active travel will depend on the duration, intensity (affected by the nature of the route, inclines and terrain), mode (e.g., walking, cycling, scooting) and frequency of the physical activity characterising the active travel^(49–56,65–79). Thus, it is important that such factors are amenable to change. The weight of the evidence from intervention and prospective studies among young people and adults combined supports the positive effects of active travel over longer periods and distances on cardiometabolic health outcomes⁽¹¹⁸⁾. Yet, further research utilising study designs that can infer causality, including acute studies, with measures of cardiometabolic disease risk other than obesity is required to draw stronger conclusions among children and adolescents specifically⁽¹¹⁸⁾. Determining the active travel characteristics that can confer health benefits would inform a health-based definition of ‘active travel’ that could be used in research and practice, which is currently lacking⁽¹¹⁸⁾.

Considering real-life application, it may be most feasible for active travel to contribute to light- or moderate-intensity physical activity; although, active commutes have potential to reach vigorous intensities in certain instances, such as up-hill cycling. It is also crucial that the duration of the commute is sufficient to contribute to daily physical activity, which may be a challenge in young children who often live in close proximity to school⁽¹¹⁹⁾. However, high levels of air pollution on some commutes may negate some of the positive health benefits, especially for children who are more susceptible than adults due to greater physiological sensitivity to air pollutants and higher respiration rates⁽¹²⁰⁾. Additionally, active travel may interact with daily diet and physical activity behaviours, and such potential compensatory responses might either negate or enhance any effects of active travel. With future research, an understanding of the acute and chronic cardiometabolic risk marker responses to active travel across different intensities, modes, durations, and frequencies has potential to inform messages that could help promote active travel for both health and climate change mitigation. Additionally, research focused on adaptation to climate change, such as strategies to keep cool during physical activity in extreme heat, will be important to promote active travel in our changing climates.

Conclusions

Cardiometabolic disease prevention is more effective than a cure and offers co-benefits for humans and the planet. Focusing on young populations to understand their distinct responses to physical activity and dietary manipulations from an early age can help inform disease prevention efforts. Knowledge of the cardiometabolic responses to acute physical activity and breakfast manipulations using controlled experimental designs among young people is growing, yet remains in its infancy when compared with the adult literature. Existing evidence suggests that acute bouts of moderate- to high-intensity exercise can acutely attenuate postprandial plasma glucose, insulin and triacylglycerol concentrations for up to 24–48-h in young people. Accumulating physical activity throughout the day with short, frequent bouts offers another potential route to attenuate postprandial cardiometabolic responses, although the benefits have not been reported consistently among young people. Likewise, although further evidence is required to draw firmer conclusions, breakfast consumption may reduce glycaemic responses to subsequent meals consumed throughout the day and enhance free-living physical activity for a possible additive health impact. If repeated habitually, such responses to physical activity and breakfast manipulations would be of benefit to long term cardiometabolic health, yet appear to depend on the study design and the physical activity, test meal and individual characteristics. To help maximise the potential cardiometabolic effects for public health, further understanding of such moderating factors and more research in samples of ‘at risk’ young people (e.g., overweight/obese, physically inactive, prediabetic) would be beneficial. Further, technological advances may offer minimally-invasive avenues to understand cellular-level responses for greater mechanistic insights. Determining whether findings based within a controlled laboratory environment translate to free-living settings with ecologically valid interventions also requires further study. Given that climate change is a serious threat that will disproportionately affect our younger generations and is relevant to cardiometabolic health, this future research should pay consideration to important planetary co-benefits.

References

1. Andes LJ, Cheng YJ, Rolka DB *et al.* (2020) Prevalence of prediabetes among adolescents and young adults in the United States, 2005-2016. *JAMA Pediatr* **174**, e194498.
2. Candler TP, Mahmoud O, Lynn RM *et al.* (2018) Continuing rise of Type 2 diabetes incidence in children and young people in the UK. *Diabet Med* **35**, 737–744.
3. Lawrence JM, Divers J, Isom S *et al.* (2021) Trends in prevalence of Type 1 and Type 2 diabetes in children and adolescents in the US, 2001-2017. *JAMA* **326**, 717–727.
4. Kit BK, Kuklina E, Carroll MD *et al.* (2015) Prevalence of and trends in dyslipidemia and blood pressure among US children and adolescents, 1999-2012. *JAMA Pediatr* **169**, 272–279.

5. Weiss R, Bremer AA & Lustig RH (2013) What is metabolic syndrome, and why are children getting it? *Ann N Y Acad Sci* **1281**, 123–140.
6. Eckelman MJ & Sherman JD (2018) Estimated Global Disease Burden From US Health Care Sector Greenhouse Gas Emissions. *Am J Public Health* **108**, Suppl. 2, S120–S122.
7. Blaak EE, Antoine JM, Benton D *et al.* (2012) Impact of postprandial glycaemia on health and prevention of disease. *Obes Rev* **13**, 923–984.
8. Heine RJ, Balkau B, Ceriello A *et al.* (2004) What does postprandial hyperglycaemia mean? *Diabet Med* **21**, 208–213.
9. Ceriello A, Esposito K, Piconi L *et al.* (2008) Oscillating glucose is more deleterious to endothelial function and oxidative stress than mean glucose in normal and type 2 diabetic patients. *Diabetes* **57**, 1349–1354.
10. Sun Y, Zhao L, Teng D *et al.* (2022) Postprandial glycemic dips are associated with metabolic disorders and CVD risk in euglycemic individuals. *J Clin Endocrinol Metab* **107**, e1631–e1642.
11. Abdul-Ghani MA, Abdul-Ghani T, Ali N *et al.* (2008) One-hour plasma glucose concentration and the metabolic syndrome identify subjects at high risk for future type 2 diabetes. *Diabetes Care* **31**, 1650–1655.
12. Barr ELM, Boyko EJ, Zimmet PZ *et al.* (2009) Continuous relationships between non-diabetic hyperglycaemia and both cardiovascular disease and all-cause mortality: Australian Diabetes, Obesity, and Lifestyle (AusDiab) study. *Diabetologia* **52**, 415–424.
13. Levitan EB, Song Y, Ford ES *et al.* (2004) Is nondiabetic hyperglycemia a risk factor for cardiovascular disease? A meta-analysis of prospective studies. *Arch Intern Med* **164**, 2147–2155.
14. Ning F, Zhang L, Dekker JM *et al.* (2012) Development of coronary heart disease and ischemic stroke in relation to fasting and 2-hour plasma glucose levels in the normal range. *Cardiovasc Diabetol* **11**, 76.
15. Cohn JS (1998) Postprandial lipemia: emerging evidence for atherogenicity of remnant lipoproteins. *Can J Cardiol* **14**, Suppl B., B18–B27.
16. Cosson E, Hamo-Tchatchouang E, Banu I *et al.* (2010) A large proportion of prediabetes and diabetes goes undiagnosed when only fasting plasma glucose and/or HbA1c are measured in overweight or obese patients. *Diabetes Metab* **36**, 312–318.
17. Goran MI & Gower BA (2001) Longitudinal study on pubertal insulin resistance. *Diabetes* **50**, 2444–2450.
18. Moran A, Jacobs DR Jr, Steinberger J *et al.* (1999) Insulin resistance during puberty: results from clamp studies in 357 children. *Diabetes* **48**, 2039–2044.

19. Jeffery AN, Metcalf BS, Hosking J *et al.* (2012) Age before stage: insulin resistance rises before the onset of puberty: a 9-year longitudinal study (EarlyBird 26). *Diabetes Care* **35**, 536–541.
20. Public Health England (2021) National Diet and Nutrition Survey: Diet and physical activity – a follow-up study during COVID-19. <https://www.gov.uk/government/statistics/ndns-diet-and-physical-activity-a-follow-up-study-during-covid-19> (accessed August 2022).
21. Kelsey MM & Zeitler PS (2016) Insulin resistance of puberty. *Curr Diab Rep* **16**, 64.
22. Roemmich JN, Clark PA, Lusk M *et al.* (2002) Pubertal alterations in growth and body composition. VI. Pubertal insulin resistance: relation to adiposity, body fat distribution and hormone release. *Int J Obes Relat Metab Disord* **26**, 701–709.
23. Cooper SB, Dring KJ, Morris JG *et al.* (2017) Sex differences in adolescents' glycaemic and insulinaemic responses to high and low glycaemic index breakfasts: a randomised control trial. *Br J Nutr* **117**, 541–547.
24. McGill HC, McMahan CA, Herderick EE *et al.* (2000) Origin of atherosclerosis in childhood and adolescence. *Am J Clin Nutr* **72**, Suppl. 5, S1307–S1315.
25. Holman RL, McGill HC, Strong JP *et al.* (1958) The natural history of atherosclerosis: the early aortic lesions as seen in New Orleans in the middle of the 20th century. *Am J Pathol* **34**, 209–235.
26. Strong JP & McGill HC (1962) The natural history of coronary atherosclerosis. *Am J Pathol* **40**, 37–49.
27. Davis PH, Dawson JD, Riley WA *et al.* (2001) Carotid intimal-medial thickness is related to cardiovascular risk factors measured from childhood through middle age: The Muscatine Study. *Circulation* **104**, 2815–2819.
28. Raitakari OT, Juonala M, Kähönen M *et al.* (2003) Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *J Am Med Assoc* **290**, 2277–2283.
29. Franks PW, Hanson RL, Knowler WC *et al.* (2010) Childhood obesity, other cardiovascular risk factors, and premature death. *N Engl J Med* **362**, 485–493.
30. Eccles JS, Midgley C, Wigfield A *et al.* (1993) Development during adolescence: The impact of stage-environment fit on young adolescents' experiences in schools and in families. *Am Psychol* **48**, 90–101.
31. Todd AS, Street SJ, Ziviani J *et al.* (2015) Overweight and obese adolescent girls: the importance of promoting sensible eating and activity behaviors from the start of the adolescent period. *Int J Environ Res Public Health* **12**, 2306–2329.

32. Henriksen EJ (2002) Invited review: Effects of acute exercise and exercise training on insulin resistance. *J Appl Physiol* **93**, 788–796.
33. Freese EC, Gist NH & Cureton KJ (2014) Effect of prior exercise on postprandial lipemia: an updated quantitative review. *J Appl Physiol* **116**, 67–75.
34. Maraki MI & Sidossis LS (2013) The latest on the effect of prior exercise on postprandial lipaemia. *Sports Med* **43**, 463–481.
35. Riddell MC (2008) The endocrine response and substrate utilization during exercise in children and adolescents. *J Appl Physiol* **105**, 725–733.
36. Timmons BW, Bar-Or O & Riddell MC (2007) Influence of age and pubertal status on substrate utilization during exercise with and without carbohydrate intake in healthy boys. *Appl Physiol Nutr Metab* **32**, 416–425.
37. Armstrong N & Welsman JR (2006) The physical activity patterns of European youth with reference to methods of assessment. *Sports Med* **36**, 1067–1086.
38. Steene-Johannessen J, Hansen BH, Dalene KE *et al.* (2020) Variations in accelerometry measured physical activity and sedentary time across Europe-harmonized analyses of 47,497 children and adolescents. *Int J Behav Nutr Phys Act* **17**, 38.
39. Guthold R, Stevens GA, Riley LM *et al.* (2020) Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1.6 million participants. *Lancet Child Adolesc Health* **4**, 23–35.
40. Farooq A, Martin A, Janssen A *et al.* (2020) Longitudinal changes in moderate-to-vigorous-intensity physical activity in children and adolescents: A systematic review and meta-analysis. *Obes Rev* **21**, e12953.
41. Monzani A, Ricotti R, Caputo M *et al.* (2019) A systematic review of the association of skipping breakfast with weight and cardiometabolic risk factors in children and adolescents. What should we better investigate in the future? *Nutrients* **11**, 387–409.
42. Betts JA, Richardson JD, Chowdhury EA *et al.* (2014) The causal role of breakfast in energy balance and health: a randomized controlled trial in lean adults. *Am J Clin Nutr* **100**, 539–547.
43. Chowdhury EA, Richardson JD, Holman GD *et al.* (2016) The causal role of breakfast in energy balance and health: a randomized controlled trial in obese adults. *Am J Clin Nutr* **103**, 747–756.
44. Yoshimura E, Hatamoto Y, Yonekura S *et al.* (2017) Skipping breakfast reduces energy intake and physical activity in healthy women who are habitual breakfast eaters: a randomized crossover trial. *Physiol Behav* **174**, 89–94

45. Chowdhury EA, Richardson JD, Tsintzas K *et al.* (2015) Carbohydrate-rich breakfast attenuates glycaemic, insulinaemic and ghrelin response to ad libitum lunch relative to morning fasting in lean adults. *Br J Nutr* **114**, 98–107.
46. Jovanovic A, Gerrard J & Taylor R (2009) The second-meal phenomenon in type 2 diabetes. *Diabetes Care* **32**, 1199–1201.
47. Jovanovic A, Leverton E, Solanky B *et al.* (2009) The second-meal phenomenon is associated with enhanced muscle glycogen storage in humans. *Clin Sci (Lond)* **117**, 119–127.
48. Gonzalez JT (2014) Paradoxical second-meal phenomenon in the acute postexercise period. *Nutrition* **30**, 961–967.
49. Zakrzewski JK & Tolfrey K (2012) Acute effect of Fatmax exercise on the metabolism in overweight and non-overweight girls. *Med Sci Sports Exerc* **44**, 1698–1705.
50. Cockcroft EJ, Williams CA, Tomlinson OW *et al.* (2015) High intensity interval exercise is an effective alternative to moderate intensity exercise for improving glucose tolerance and insulin sensitivity in adolescent boys. *J Sci Med Sport* **18**, 720–724.
51. Cockcroft EJ, Williams CA, Weaver H *et al.* (2017) Acute exercise and insulin sensitivity in boys: a time-course study. *Int J Sports Med* **38**, 967–974.
52. Cockcroft EJ, Williams CA, Jackman SR *et al.* (2018) A single bout of high-intensity interval exercise and work-matched moderate-intensity exercise has minimal effect on glucose tolerance and insulin sensitivity in 7- to 10-year-old boys. *J Sports Sci* **36**, 149–155.
53. Dring KJ, Cooper SB, Morris JG *et al.* (2019) Cytokine, glycemic, and insulinemic responses to an acute bout of games-based activity in adolescents. *Scand J Med Sci Sports* **29**, 597–605.
54. Williams RA, Cooper S, Dring KJ *et al.* (2021) Effect of acute football activity and physical fitness on glycaemic and insulinaemic responses in adolescents. *J Sports Sci* **39**, 1127–1135.
55. Short KR, Pratt LV, Teague AM *et al.* (2013) Postprandial improvement in insulin sensitivity after a single exercise session in adolescents with low aerobic fitness and physical activity. *Pediatr Diabetes* **14**, 129–137.
56. Dring KJ, Cooper SB, Williams RA *et al.* (2020) Effect of exercise duration on postprandial glycaemic and insulinaemic responses in adolescents. *Nutrients* **12**, 754–764.
57. Holloway GP, Bonen A & Spriet LL (2009) Regulation of skeletal muscle mitochondrial fatty acid metabolism in lean and obese individuals. *Am J Clin Nutr* **89**, S455–S462.

58. Schenk S & Horowitz JF (2007) Acute exercise increases triglyceride synthesis in skeletal muscle and prevents fatty acid–induced insulin resistance. *J Clin Invest* **117**, 1690–1698.
59. Sinha R, Dufour S, Petersen KF *et al.* (2002) Assessment of skeletal muscle triglyceride content by (1)H nuclear magnetic resonance spectroscopy in lean and obese adolescents: relationships to insulin sensitivity, total body fat, and central adiposity. *Diabetes* **51**, 1022–1027.
60. Eriksson BO, Gollnick PD & Saltin B (1973) Muscle metabolism and enzyme activities after training in boys 11–13 years old. *Acta Physiol Scand* **87**, 485–499.
61. Eriksson BO (1980) Muscle metabolism in children—a review. *Acta Physiol Scand* **283**, 20–28.
62. Bond B, Williams CA, Isic C *et al.* (2015) Exercise intensity and postprandial health outcomes in adolescents. *Eur J Appl Physiol* **115**, 927–936.
63. Bond B, Gates PE, Jackman SR *et al.* (2015) Exercise intensity and the protection from postprandial vascular dysfunction in adolescents. *Am J Physiol Heart Circ Physiol* **308**, H1443–H1450.
64. Ghafouri K, Cooney J, Bedford DK *et al.* (2015) Moderate exercise increases affinity of large very low-density lipoproteins for hydrolysis by lipoprotein lipase. *J Clin Endocrinol Metab* **100**, 2205–2213.
65. Tolfrey K, Doggett A, Boyd C *et al.* (2008) Postprandial triacylglycerol in adolescent boys: a case for moderate exercise. *Med Sci Sports Exerc* **40**, 1049–1057.
66. Tolfrey K, Bentley C, Goad M *et al.* (2012) Effect of energy expenditure on postprandial triacylglycerol in adolescent boys. *Eur J Appl Physiol* **112**, 23–31.
67. Tolfrey K, Engstrom A, Murphy C *et al.* (2014) Exercise energy expenditure and postprandial lipaemia in girls. *Med Sci Sports Exerc* **46**, 239–246.
68. Barrett LA, Morris JG, Stensel DJ *et al.* (2007) Exercise and postprandial plasma triacylglycerol concentrations in healthy adolescent boys. *Med Sci Sports Exerc* **39**, 116–122.
69. MacEneaney OJ, Harrison M, O’Gorman DJ *et al.* (2009) Effect of prior exercise on postprandial lipemia and markers of inflammation and endothelial activation in normal weight and overweight adolescent boys. *Eur J Appl Physiol* **106**, 721–729.
70. Lee S, Burns SF, White D *et al.* (2013) Effects of acute exercise on postprandial triglyceride response after a high-fat meal in overweight black and white adolescents. *Int J Obes* **37**, 966–971.
71. Sedgwick MJ, Morris JG, Nevill ME *et al.* (2013) Effect of exercise on postprandial endothelial function in adolescent boys. *Br J Nutr* **110**, 301–309.

72. Sedgwick MJ, Morris JG, Nevill ME *et al.* (2014) The accumulation of exercise and postprandial endothelial function in boys. *Scand J Med Sci Sports* **24**, e11–e19.
73. Thackray AE, Barrett LA & Tolfrey K (2015) Acute effects of energy deficit induced by moderate-intensity exercise or energy-intake restriction on postprandial lipemia in healthy girls. *Pediatr Exerc Sci* **27**, 192–202.
74. Thackray AE, Barrett LA & Tolfrey K (2016) Energy replacement diminishes the effect of exercise on postprandial lipemia in boys. *Metabolism* **65**, 496–506.
75. Smallcombe JW, Barrett LA, Morris JG *et al.* (2018) Small-sided soccer in school reduces postprandial lipemia in adolescent boys. *Med Sci Sports Exerc* **50**, 2351–2359.
76. Thackray AE, Barrett LA & Tolfrey K (2013) Acute high-intensity interval running reduces postprandial lipemia in boys. *Med Sci Sports Exerc* **45**, 1277–1284.
77. Thackray AE, Barrett LA & Tolfrey K (2016) High-intensity running and energy restriction reduces postprandial lipemia in girls. *Med Sci Sports Exerc* **48**, 402–411.
78. Smallcombe JW, Barrett LA, Sherar LB *et al.* (2020) Short sprints accumulated at school modulate postprandial metabolism in boys. *Med Sci Sports Exerc* **52**, 67–76.
79. Sedgwick MJ, Morris JG, Nevill ME *et al.* (2015) Effect of repeated sprints on postprandial endothelial function and triacylglycerol concentrations in adolescent boys. *J Sports Sci* **33**, 806–816.
80. Bauman AE, Petersen CB, Blond K *et al.* (2018) The descriptive epidemiology of sedentary behaviour. In *Sedentary Behaviour Epidemiology. Springer Series on Epidemiology and Public Health.*, pp. 73–106 [MF Leitzmann, C Jochem and D Schmid, editors]. Springer, Cham.
81. Tremblay MS, Aubert S, Barnes JD *et al.* (2017) Sedentary Behavior Research Network (SBRN) - Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act* **14**, 75-91.
82. Ekelund U, Tarp J, Steene-Johannessen J *et al.* (2019) Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ* **366**, 14570.
83. Patterson R, McNamara E, Tainio M *et al.* (2018) Sedentary behaviour and risk of all-cause, cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review and dose response meta-analysis. *Eur J Epidemiol* **33**, 811–829.
84. Carson V, Hunter S, Kuzik N *et al.* (2016) Systematic review of sedentary behaviour and health indicators in school-aged children and youth: an update. *Appl Physiol Nutr Metab* **41**, S240–S265.

85. Timmons BW & Bar-Or O (2003) RPE during prolonged cycling with and without carbohydrate ingestion in boys and men. *Med Sci Sports Exerc* **35**, 1901–1907.
86. Saunders TJ, Atkinson HF, Burr J *et al.* (2018) The acute metabolic and vascular impact of interrupting prolonged sitting: a systematic review and meta-analysis. *Sports Med* **48**, 2347–2366.
87. Belcher BR, Berrigan D, Papachristopoulou A *et al.* (2015) Effects of interrupting children's sedentary behaviors with activity on metabolic function: A randomized trial. *J Clin Endocrinol Metab* **100**, 3735–3743.
88. Broadney MM, Belcher BR, Berrigan DA *et al.* (2018) Effects of interrupting sedentary behavior with short bouts of moderate physical activity on glucose tolerance in children with overweight and obesity: A randomized crossover trial. *Diabetes Care* **41**, 2220–2228.
89. Bond B, Williams CA, Jackman SR *et al.* (2015) Accumulating exercise and postprandial health in adolescents. *Metabolism* **64**, 1068–1076.
90. Saunders TJ, Chaput JP, Goldfield GS *et al.* (2013) Prolonged sitting and markers of cardiometabolic disease risk in children and youth: a randomized crossover study. *Metabolism* **62**, 1423–1428.
91. Smallcombe JW, Biddle GJH, Slater T *et al.* (2022) Breaking sitting time with physical activity increases energy expenditure but does not alter postprandial metabolism in girls. *Med Sci Sports Exerc.* Published online: 17 June 2022. doi: 10.1249/MSS.0000000000002979.
92. Fletcher EA, Salmon J, McNaughton SA *et al.* (2018) Effects of breaking up sitting on adolescents' postprandial glucose after consuming meals varying in energy: a cross-over randomised trial. *J Sci Med Sport* **21**, 280–285.
93. Souza MR, Neves MEA, Gorgulho BM *et al.* (2021) Breakfast skipping and cardiometabolic risk factors in adolescents: Systematic review. *Rev Saude Publica* **55**, 107–128.
94. Ballon A, Neuenschwander M & Schlesinger S (2019) Breakfast skipping is associated with increased risk of type 2 diabetes among adults: A systematic review and meta-analysis of prospective cohort studies *J Nutr* **149**, 106–113.
95. Chen H, Zhang B, Ge Y *et al.* (2020) Association between skipping breakfast and risk of cardiovascular disease and all cause mortality: A meta-analysis. *Clin Nutr* **39**, 2982–2988.
96. Harrell JS, McMurray RG, Baggett CD *et al.* (2005) Energy costs of physical activities in children and adolescents. *Med Sci Sports Exerc* **37**, 329–336.

97. Rogol AD, Clark PA & Roemmich JN (2000) Growth and pubertal development in children and adolescents: Effects of diet and physical activity. *Am J Clin Nutr* **72**, Suppl. 2, S521–S528.
98. Corder K, van Sluijs EM, Steele RM *et al.* (2011) Breakfast consumption and physical activity in British adolescents. *Br J Nutr* **105**, 316–321.
99. Corder K, van Sluijs EM, Ridgway CL *et al.* (2014) Breakfast consumption and physical activity in adolescents: daily associations and hourly patterns. *Am J Clin Nutr* **99**, 361–368.
100. Vissers PA, Jones AP, Corder K *et al.* (2013) Breakfast consumption and daily physical activity in 9-10-year-old British children. *Public Health Nutr* **16**, 1281–1290.
101. Zakrzewski-Fruer JK, Plekhanova T, Mandila D *et al.* (2017) Effect of breakfast omission and consumption on energy intake and physical activity in adolescent girls: A randomised controlled trial. *Br J Nutr* **118**, 392–400.
102. Zakrzewski-Fruer JK, Wells EK, Crawford NSG *et al.* (2018) Physical activity duration but not energy expenditure differs between daily compared with intermittent breakfast consumption in adolescent girls: a randomized crossover trial. *J Nutr* **148**, 236–244.
103. Zakrzewski-Fruer JK, Seall C & Tolfrey K (2021) Breakfast consumption suppresses appetite but does not increase daily energy intake or physical activity energy expenditure when compared with breakfast omission in adolescent girls who habitually skip breakfast: a 7-day randomised crossover trial. *Nutrients* **13**, 4261–4278.
104. O’Neil CE, Byrd-Bredbenner C, Hayes D *et al.* (2014) The role of breakfast in health: Definition and criteria for a quality breakfast. *J Acad Nutr Diet* **114**, Suppl. 12, S8–S26.
105. Eriksson BO & Saltin B (1974) Muscle metabolism during exercise in boys aged 11 to 16 years compared to adults. *Acta Paediatr Belg* **28**, 257–265.
106. Kovacic K, Zhang L, Nugent Liegl M *et al.* (2021) Gastric emptying in healthy children using the Spirulina breath test: The impact of gender, body size, and pubertal development. *Neurogastroenterol Motil* **33**, e14063.
107. Alwattar AY, Thyfault JP & Leidy HJ (2015) The effect of breakfast type and frequency of consumption on glycemic response in overweight/obese late adolescent girls. *Eur J Clin Nutr* **69**, 885–90.
108. Bauer LB, Reynolds LJ, Douglas SM *et al.* (2015) A pilot study examining the effects of consuming a high-protein vs normal-protein breakfast on free-living glycemic control in overweight/obese 'breakfast skipping' adolescents. *Int J Obes (Lond)* **39**, 1421–1424.
109. Leidy HJ, Hoertel HA, Douglas SM *et al.* (2015) A high-protein breakfast prevents body fat gain, through reductions in daily intake and hunger, in ‘Breakfast skipping’ adolescents. *Obesity (Silver Spring)* **23**, 1761–1764.

110. Bonuccelli S, Muscelli E, Gastaldelli A *et al.* (2009) Improved tolerance to sequential glucose loading (Staub-Traugott effect): size and mechanisms. *Am J Physiol Endocrinol Metab* **297**, e532–e537.
111. Liljeberg H & Björck I (2000) Effects of a low-glycaemic index spaghetti meal on glucose tolerance and lipaemia at a subsequent meal in healthy subjects. *Eur J Clin Nutr* **54**, 24–28.
112. Gibney MJ, Barr SI, Bellisle F *et al.* (2018) Towards an evidence-based recommendation for a balanced breakfast-A proposal from the International Breakfast Research Initiative. *Nutrients* **10**, 1540–1555.
113. Monteagudo C, Palacin-Arce A, Mdel MB *et al.* (2013) Proposal for a Breakfast Quality Index (BQI) for children and adolescents. *Public Health Nutr* **16**, 639–644.
114. Zakrzewski JK, Stevenson EJ & Tolfrey K (2012) Effect of breakfast glycaemic index on metabolic responses during rest and exercise in overweight and non-overweight adolescent girls. *Eur J Clin Nutr* **66**, 436–442.
115. Zakrzewski-Fruer JK, Morari V, Champion RB *et al.* (2021) Acute cardiometabolic and exercise responses to breakfast consumption versus breakfast omission in adolescent girls: A crossover trial. Abstracts from the 32nd Pediatric Work Physiology Conference (September 2021, Virtual). *Pediatr Exerc Sci* **33**, Suppl 1., S1-S21.
116. Gasparri G, Eissa M, Imbago-Jácome D *et al.* (2022) Climate emergency: how should COP27 do better for adolescents and young people? *BMJ* **376**, o816.
117. Martins J, Costa J, Sarmiento H *et al.* (2021) Adolescents' perspectives on the barriers and facilitators of physical activity: An updated systematic review of qualitative studies. *Int J Environ Res Public Health* **18**, 4954–4978.
118. Saunders LE, Green JM, Petticrew MP *et al.* (2013) What are the health benefits of active travel? A systematic review of trials and cohort studies. *PLoS One* **8**, e69912.
119. Metcalf B, Voss L, Jeffery A *et al.* (2004) Physical activity cost of the school run: impact on schoolchildren of being driven to school (EarlyBird22). *BMJ* **329**, 832–833.
120. Al-Kindi SG, Brook RD, Biswal S *et al.* (2020) Environmental determinants of cardiovascular disease: lessons learned from air pollution. *Nat Rev Cardiol* **17**, 656–672.

Figure legends

Fig. 1. Potential avenues that support a focus on children and adolescents (young people) for cardiometabolic disease prevention through physical activity and/or diet with co-benefits to mitigating climate change. Note that examples of pathways linking physical activity and dietary

manipulations in young people with climate change mitigation include, but are not limited to, active travel, human-nature connections and plant-based diets.

Fig. 2. (A) Schematic of typical protocols adopted to assess the postprandial cardiometabolic responses to acute exercise or physical activity bouts. Typically, a 1- or 2-day protocol is used, with exercise/physical activity manipulations and rest (control) conditions performed on day 1 and postprandial cardiometabolic risk marker responses assessed on day 1 and/or day 2 and in some cases day 3. Protocols differ according to whether the primary outcome is plasma glucose and insulin or triacylglycerol. Repeated capillary or venous blood samples are collected at pre-determined time-points in the fasted and postprandial state. OGTT: oral glucose tolerance test; HCM: high-carbohydrate meal; HFM: high-fat meal.

(B) Illustration of the typical circulating plasma glucose and triacylglycerol response to consumption of a glucose load or meal at 0 h (indicated by the arrow). Postprandial glycaemic and insulinaemic responses are typically immediate, peaking at ~15–30-min and returning to baseline within ~2-h^(20,49,56,114), whereas postprandial triacylglycerol rises slower to reach a peak at ~3–4-h before returning to baseline after 6–8-h in the absence of a second meal challenge^(65–68). As such, glucose and insulin concentrations require more frequent sampling over a shorter duration often in response to a carbohydrate-based test meal/drink, whereas less frequent blood sampling over a longer duration is required to capture postprandial triacylglycerol profiles, often in response to a high-fat meal. The macronutrient content of the test meal is typically designed to induce a sufficient metabolic challenge specific to the primary cardiometabolic risk marker under investigation to ensure ample scope for intervention effects. Mixed macronutrient meals have also been employed for the measurement of both glycaemia and lipaemia or to enhance ecological validity.

Fig. 1.




Fig. 2.

